



Published in final edited form as:

*J Vasc Surg.* 2024 November ; 80(5): 1455–1463. doi:10.1016/j.jvs.2024.05.048.

## Seven Years of The Transcarotid Artery Revascularization Surveillance Project, Comparison To Transfemoral Stenting And Endarterectomy

Sabrina Straus, BS<sup>1,2</sup>, Sai Divya Yadavalli, MD<sup>1</sup>, Sara Allievi, MD<sup>1,3</sup>, Andrew Sanders, MD<sup>1</sup>, Roger B. Davis, ScD<sup>1</sup>, Mahmoud B Malas, MD<sup>2</sup>, Grace J Wang, MD<sup>4</sup>, Vikram S Kashyap, MD<sup>5</sup>, Jack Cronenwett, MD<sup>6</sup>, Raghu L Motaganahalli, MD<sup>7</sup>, Brian Nolan, MD<sup>8</sup>, Jens Eldrup-Jorgensen, MD<sup>8</sup>, Marc Schermerhorn, MD<sup>1</sup>

<sup>1</sup>Department of Surgery, Division of Vascular and Endovascular Surgery, Beth Israel Deaconess Medical Center, Harvard Medical School, Boston, MA, USA.

<sup>2</sup>Department of Surgery, Division of Vascular and Endovascular Surgery, UC San Diego (UCSD), San Diego, California

<sup>3</sup>Department of Vascular Surgery, Fondazione IRCCS Cà Granda Ospedale Maggiore Policlinico, Milan, Italy

<sup>4</sup>Division of Vascular Surgery and Endovascular Therapy, Department of Surgery, Hospital of the University of Pennsylvania, Philadelphia, PA.

<sup>5</sup>Frederik Meijer Heart and Vascular Institute, Corewell Health, Grand Rapids, MI

<sup>6</sup>Section of Vascular Surgery, Dartmouth Medical School, 1 Medical Center Drive, Lebanon, NH.

<sup>7</sup>Division of Vascular Surgery, Department of Surgery, Indiana University School of Medicine, Indianapolis, IN.

<sup>8</sup>Division of Vascular and Endovascular Therapy, Department of Surgery, Maine Medical Center, Portland, ME.

### Abstract

**Objective:** This study utilizes the latest data from the Vascular Quality Initiative (VQI), which now encompasses over 50,000 transcarotid artery revascularization (TCAR) procedures, to offer a sizeable dataset for comparing the effectiveness and safety of TCAR, transfemoral carotid artery stenting (tfCAS), and carotid endarterectomy (CEA). Given this substantial dataset, we are

---

**Corresponding author:** Marc L. Schermerhorn, MD, FACS, Beth Israel Deaconess Medical Center, 110 Francis Street, Suite 5B, Boston, MA 02215, mscherm@bidmc.harvard.edu.

**Publisher's Disclaimer:** This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

Conflicts of Interest  
None

Presentation information

Accepted for Poster Competition at Vascular Annual Meeting of the Society for Vascular Surgery, Chicago, IL, June 19, 2024

now able to compare outcomes overall and stratified by symptom status across revascularization techniques.

**Methods:** Utilizing VQI data from September 2016 to August 2023, we conducted a risk-adjusted analysis by applying inverse probability of treatment weighting to compare in-hospital outcomes between TCAR vs tfCAS, CEA vs tfCAS, and TCAR vs CEA. Our primary outcome measure was in-hospital stroke/death. Secondary outcomes included myocardial infarction and cranial nerve injury.

**Results:** A total of 50,068 patients underwent TCAR, 25,361 patients underwent tfCAS, and 122,737 patients underwent CEA. TCAR patients were older, more likely to have coronary artery disease, chronic kidney disease, and undergo coronary artery bypass grafting/percutaneous coronary intervention as well as prior contralateral CEA/CAS compared to both CEA and tfCAS. TfCAS had higher odds of stroke/death when compared with TCAR (2.9% vs 1.6%, aOR=1.84, 95% CI:1.65–2.06; P<.001) and CEA (2.9% vs 1.3%, aOR=2.21, 95% CI:2.01–2.43; P<.001). CEA had slightly lower odds of stroke/death compared with TCAR (1.3% vs 1.6%, aOR=0.83, 95% CI:0.76–0.91; P<.001). TfCAS had lower odds of cranial nerve injury compared with TCAR (0.0% vs 0.3%, aOR=0.00, 95% CI:0.00–0.00; P<.001) and CEA (0.0% vs 2.3%, aOR=0.00, 95% CI:0.0–0.0; P<.001) as well as lower odds of myocardial infarction compared with CEA (0.4% vs 0.6%, aOR=0.67, 95% CI:0.54–0.84; P<.001). CEA compared with TCAR had higher odds of myocardial infarction (0.6% vs 0.5%, aOR=1.31, 95% CI:1.13–1.54; P<.001) and cranial nerve injury (2.3% vs 0.3%, aOR=9.42, 95% CI:7.78–11.4; P<.001).

**Conclusions:** While tfCAS may be beneficial for select patients, the lower stroke/death rates associated with CEA and TCAR are preferred. When deciding between CEA and TCAR, it's important to weigh additional procedural factors and outcomes such as myocardial infarction and cranial nerve injury, particularly when stroke/death rates are similar. Additionally, evaluating subgroups that may benefit from one procedure over another is essential for informed decision-making and enhanced patient care in the treatment of carotid stenosis.

### Table of Contents Summary:

In this retrospective review of over 50,000 transcarotid artery revascularization, 25,000 transfemoral carotid artery stenting, and 120,000 carotid endarterectomy patients, transfemoral carotid artery stenting had the highest rates of stroke/death. Given the relatively smaller differences in stroke/death rates between transcarotid artery revascularization and endarterectomy, evaluation of additional outcomes is important.

### Keywords

Carotid revascularization; transcarotid artery revascularization; transfemoral carotid artery stenting; carotid endarterectomy

### Introduction

Among the various approaches for carotid revascularization, carotid endarterectomy (CEA) and transfemoral carotid artery stenting (tfCAS) are the most established options.<sup>1</sup> While

CEA is the standard treatment, tfCAS was introduced as an alternative in high-risk patients with high-grade asymptomatic (>70%) or symptomatic stenosis.<sup>2, 3</sup>

After FDA approval in 2015,<sup>4</sup> transcarotid artery revascularization (TCAR) emerged as a potential alternative, providing cerebral protection through its flow reversal mechanism and eliminating the need for aortic arch crossing.<sup>5, 6</sup> Several retrospective studies have found TCAR consistently outperforms tfCAS, with lower rates of stroke/death during in-hospital stay and at 1-year follow-up.<sup>3, 8</sup> When comparing stroke/death outcomes between TCAR and CEA, studies have found no significant differences; however, when stratifying by symptomatic status, studies present conflicting results regarding outcomes.<sup>7, 8, 9, 10</sup> Nevertheless, TCAR's low reported stroke/death rates have led to its expanded indications and adoption by a considerable portion of physicians.<sup>11, 12, 13, 14, 15</sup>

When comparing CEA with tfCAS, higher in-hospital, 30-day, and long-term stroke/death rates have been observed for tfCAS in symptomatic patients, with no statistical difference in asymptomatic patients in retrospective studies and meta-analyses.<sup>16, 17, 18, 19</sup> The findings from Carotid Revascularization Endarterectomy vs. Stenting Trial (CREST) align with these observations, attributing the lack of significance in the asymptomatic subanalysis to lower statistical power.<sup>20</sup> The insufficient statistical power and lack of consensus emphasize the necessity for a thorough evaluation of in-hospital outcomes across all three procedures. Now with over 50,000 TCAR patients in the VQI dataset, there is a substantial sample size to provide updated insights into the comparative safety of TCAR, tfCAS, and CEA and stratify by symptom status.

## Methods

### Data Source

This is a retrospective cohort study using the Society for Vascular Surgery Vascular Quality Initiative (SVS-VQI) registry database ([www.vqi.org](http://www.vqi.org)). The VQI database includes information from over 1,000 international participating medical centers and provides de-identified data on major vascular procedures. Importantly, the TCAR Surveillance Project mandates that institutions conducting TCAR procedures adhere to the requirement of entering all TCAR and CAS data into the VQI registry to be eligible for Medicare reimbursement. The STROBE guidelines (<https://www.strobe-statement.org/>) were used to ensure proper reporting of methods, results, and discussion. Ethical approval for this study was provided by the VQI Research Advisory Committee and the Institutional Review Board at the Beth Israel Deaconess Medical Center. The need for informed consent was waived due to the retrospective and de-identified nature of the data.

### Patient Cohort

From September 2016 to August 2023, all patients undergoing CAS (n=92,217), including both TCAR and tfCAS, and CEA (n=126,582) in the VQI were identified, aligning with the commencement of the TCAR Surveillance Project. Patients under 18 years-old were excluded. In the CAS cohort, patients with non-atherosclerotic lesions (i.e., trauma, dissection, and fibromuscular dysplasia) (n=3,438), more than one treated lesion (n=1,019),

tandem internal carotid artery lesion >70% stenosis (n=3,132) and planned intracerebral intervention (n=1,498) were excluded. In the CEA cohort, patients undergoing other concomitant endovascular procedures, CABG, or other arterial interventions were excluded (n=3,607). Records with missing information regarding procedure type (TCAR, tfCAS, CEA) or symptom status were excluded.

### Variable definitions and outcomes

The estimated glomerular filtration rate (eGFR) was calculated using the new Chronic Kidney Disease Epidemiology (CKD-EPI) Creatinine Equation (2021).<sup>21</sup> Patients were grouped based on eGFR as follows: eGFR>45 mL/min/1.73m<sup>2</sup>, eGFR 30–45 mL/min/1.73m<sup>2</sup>, eGFR<30 mL/min/1.73m<sup>2</sup>, or preoperative dialysis (regardless of eGFR). Preoperative anemia was defined as hemoglobin <10 g/dL.<sup>22</sup> Symptom status was defined as the presence of ipsilateral ocular, ipsilateral cortical TIA, or ipsilateral stroke symptoms documented in medical records within six months prior to the index procedure in accordance with SVS reporting standards.<sup>23</sup>

The primary outcome was the composite variable in-hospital stroke/death. Secondary outcomes included in-hospital stroke, death, myocardial infarction (MI), composite variable stroke/death/MI, cranial nerve injury (CNI), prolonged length of stay (≥ 2 days),<sup>24</sup> and bleeding. Stroke, MI, CNI, and bleeding were all defined according to the VQI criteria. Stroke was defined as the presence of symptoms lasting 24 hours or longer, involving neurological or visual deficits. MI was indicated by troponin level, EKG, or diagnosed clinically according to American College of Cardiology criteria.<sup>25</sup> CNI was defined as any occurrence of cranial nerve injuries (CN7, CN9, CN10, CN12) that began after the procedure. Bleeding was defined as procedure-related bleeding resulting in surgical repair or intervention, such as percutaneous thrombin injection or covered stent placement.

### Statistical Analysis

We compared demographics, comorbidities, and procedural characteristics across the three procedures (TCAR, tfCAS, CEA). Categorical variables were presented as counts and percentages and were compared using Pearson's Chi-square or Fisher's exact tests as appropriate. Continuous variables were presented as mean ± standard deviation and compared using Student's t-test or median IQR. For non-normal distribution, the Wilcoxon rank sum test was utilized.

To account for differences among the three groups of patients, we used inverse probability of treatment weighting (IPTW) based on propensity scores.<sup>26, 27</sup> In contrast to pairwise propensity score matching, IPTW offers the advantage of leveraging the entire sample without constraints on the covariates included in the model. By accounting for covariates related to treatment selection, such as background demographics, comorbidities, symptom status, procedure history, and surgery year, IPTW ensures thorough adjustment and efficient utilization of all available data.<sup>28</sup> The covariates used were age, sex, race, BMI, hypertension, diabetes, coronary artery disease (CAD), prior congestive heart failure (CHF), preoperative smoking, chronic obstructive pulmonary disease (COPD), renal dysfunction, anemia, symptomatic amaurosis, symptomatic hemispheric TIA, symptomatic stroke, prior

Coronary Artery Bypass Grafting (CABG)/Percutaneous Coronary Intervention (PCI), prior contralateral CEA/CAS, % stenosis of ipsilateral and contralateral carotid, and surgery year. Sensitivity analysis tested for differences after trimming extreme variables, but no differences were noted. We compared the weighted treatment groups in terms of baseline characteristics to assess the success of the propensity score weighting in achieving balance across the groups. Standardized mean differences (SMDs) were calculated for each covariate, and an SMD <.10 was considered indicative of a balanced distribution. Tests that compared all 3 groups simultaneously by using chi-squared tests involving higher degrees-of-freedom and F-tests for continuous predictors were conducted as appropriate. The in-hospital outcomes were reported in terms of adjusted rates and odds ratios for each of the treatment comparisons. A post-hoc analysis was also conducted to confirm that this study was sufficiently powered. As a subanalysis, we evaluated in-hospital outcomes, stratifying by symptom status.

Stata MP version 17.0 was used to combine datasets, define variables, and stratify data by carotid revascularization methodology. All subsequent statistical analyses were performed using R version 4.3.1 (<http://www.r-project.org>). All tests were two-sided and p-values <.05 were considered significant.

## Results

### All Patients

After exclusions, a total of 50,068 (25%) patients underwent TCAR, 25,361(13%) patients underwent tfCAS, and 122,737 (62%) patients underwent CEA during the study period (Table I). Before IPTW, TCAR patients compared with tfCAS and CEA were older, had more CAD and CKD, and were more likely to undergo CABG/PCI as well as prior contralateral CEA/CAS (Table I). Patients who received tfCAS were more likely to undergo local anesthesia (81% vs 13% vs 6%;  $P<.001$ ), had minimal protamine usage (12% vs 88% vs 75%;  $P<.001$ ), and experienced the shortest procedure duration ( $67\pm 40$  vs  $70\pm 55$  vs  $117\pm 45$ ;  $P<.001$ ) compared to TCAR and CEA respectively.

Adjusted rates for baseline characteristics after IPTW were included (Table I). In-hospital outcomes were reported for the overall patient population (Table II), as well as for symptomatic (Table III) and asymptomatic patients (Table IV), comparing TCAR vs tfCAS, CEA vs tfCAS, and TCAR vs CEA. Adjusted stroke/death rates per year by procedure type were also plotted (Figure 1).

Prior to adjustment, the in-hospital stroke/death rates were highest for tfCAS compared to TCAR and CEA (2.8% vs 1.5% vs 1.3%). After adjustment (Table II), tfCAS compared with TCAR had higher odds of stroke/death (2.9% vs 1.6%; aOR=1.84, 95% CI:1.65–2.06;  $P<.001$ ). For secondary outcomes, tfCAS also had higher odds of stroke (2.1% vs 1.3%; aOR=1.58, 95% CI:1.39–1.79;  $P<.001$ ), death (1.1% vs 0.4%, aOR=2.80, 95% CI:2.29–3.42;  $P<.001$ ), stroke/death/MI (3.2% vs 2.0%, aOR=1.62, 95% CI:1.46–1.80;  $P<.001$ ), and prolonged length of stay (35.5% vs 29.6%; aOR=1.31, 95% CI:1.27–1.36;  $P<.001$ ) compared to TCAR. However, tfCAS had lower odds of CNI (0.0% vs 0.3%; aOR=0.00,

95% CI:0.00–0.00;  $P<.001$ ) and bleeding (0.4% vs 0.8%; aOR=0.52, 95% CI:0.41–0.67;  $P<.001$ ).

Similarly, tfCAS compared with CEA had higher odds of stroke/death (2.9% vs 1.3%; aOR=2.21, 95% CI:2.01–2.43;  $P<.001$ ) as well as secondary outcomes: stroke, death, stroke/death/MI, and prolonged length of stay (Table II). However, tfCAS had lower odds of in-hospital MI (0.4% vs 0.6%; aOR=0.67, 95% CI:0.54–0.84;  $P<.001$ ), CNI (0.0% vs 2.3%; aOR=0.00, 95% CI:0.00–0.00;  $P<.001$ ), and bleeding (0.4% vs 1.0%; aOR=0.38, 95% CI:0.30–0.47;  $P<.001$ ).

On the other hand, CEA compared with TCAR had lower odds of stroke/death (1.3% vs 1.6%; aOR=0.83, 95% CI:0.76–0.91;  $P<.001$ ), stroke (1.1% vs 1.3%; aOR=0.84, 95% CI:0.76–0.93;  $P<.001$ ), and death (0.3% vs 0.4%, aOR=0.76, 95% CI:0.64–0.92;  $P=.004$ ), similar odds of stroke/death/MI (1.9% vs 2.0%, aOR=0.94, 95% CI:0.87–1.02;  $P=.14$ ), and higher odds of MI (0.6% vs 0.5%, aOR=1.31, 95% CI:1.13–1.54;  $P<.001$ ), prolonged length of stay (30.1% vs 29.6%; aOR=1.02, 95% CI:1.00–1.05;  $P=.047$ ), CNI (2.3% vs 0.3%; aOR=9.42, 95% CI:7.78–11.4;  $P<.001$ ), and bleeding (1.0% vs 0.8%; aOR=1.38, 95% CI:1.22–1.57;  $P<.001$ ).

### Symptomatic Patients

TfCAS compared with TCAR still had higher odds of stroke/death (4.4% vs 2.4%; aOR=1.87, 95% CI:1.59–2.21;  $P<.001$ ) and notably death (1.9% vs 0.6%; aOR=3.35, 95% CI:2.50–4.48;  $P<.001$ ) and prolonged length of stay (52.2% vs 41.1%; aOR=1.56, 95% CI:1.47–1.66;  $P<.001$ ) (Table III). Similarly, tfCAS compared with CEA still had higher odds of stroke/death (4.4% vs 2.1%; aOR=2.18, 95% CI:1.91–2.48;  $P<.001$ ) and prolonged length of stay (52.2% vs 42.9%; aOR=1.45, 95% CI:1.38–1.53;  $P<.001$ ). Nevertheless, tfCAS still had the lowest rates of CNI compared with both TCAR and CEA.

CEA compared with TCAR still had lower odds of stroke/death (2.1% vs 2.4%; aOR=0.86, 95% CI:0.75–0.99;  $P=.039$ ) and stroke (1.8% vs 2.1%; aOR=0.84, 95% CI:0.72–0.98;  $P=.029$ ), although death was no longer significant (0.5% vs 0.6%; aOR=0.83, 95% CI:0.62–1.10;  $P=.2$ ). Furthermore, CEA still had higher odds of in-hospital MI (0.6% vs 0.4%; aOR=1.44, 95% CI:1.05–1.96;  $P=.022$ ), prolonged length of stay (42.9% vs 41.1%; aOR=1.08, 95% CI:1.03–1.12;  $P<.001$ ), CNI (2.8% vs 0.3%; aOR=8.60, 95% CI:6.20–11.9;  $P<.001$ ).

### Asymptomatic Patients

TfCAS still had higher odds of stroke/death compared with TCAR (2.1% vs 1.3%; aOR=1.68, 95% CI:1.44–1.96;  $P<.001$ ) and CEA (2.1% vs 1.0%; aOR=2.15, 95% CI:1.87–2.47;  $P<.001$ ). When comparing tfCAS with TCAR, tfCAS patients had similar although now statistically significant lower odds of MI (0.3% vs 0.5%; aOR=0.67, 95% CI:0.48–0.94;  $P=.020$ ). CEA compared with TCAR still had lower odds of stroke/death (1.0% vs 1.3%; aOR=0.78, 95% CI:0.70–0.88;  $P<.001$ ). The remaining outcomes remained similar to the overall unstratified population (Table IV).

## Discussion

By utilizing a modern database with almost 200,000 carotid revascularization procedures, our study reveals important, updated findings regarding the comparative safety of carotid revascularization techniques. In the overall population, rates of stroke and stroke/death were statistically lowest for CEA compared to TCAR and tfCAS. When comparing the difference in stroke/death rates between the three groups, the difference in stroke/death rates between TCAR and CEA (0.3%) is relatively smaller compared to that of tfCAS to TCAR (1.3%) or tfCAS to CEA (1.6%). As a result, when comparing CEA and TCAR, additional outcomes such as MI and CNI should factor into clinical decision-making. Particularly, when evaluating CNI and length of stay, TCAR consistently exhibited superior outcomes compared with CEA. As a result, the choice between TCAR versus CEA should be based on patient risk factors and anatomy, using shared decision making. On the other hand, the relatively larger discrepancies in stroke/death outcomes associated with tfCAS may suggest that tfCAS should be reserved for select circumstances, such as prohibitive neck anatomy.<sup>29</sup>

When comparing TCAR with CEA, previous literature has found that TCAR performs similar to CEA in terms of post-operative outcomes for all patients and when stratifying by symptomatic status, consistent with our current findings.<sup>9, 10</sup> In previous studies comparing in-hospital outcomes between TCAR and CEA using VQI data, both analyses found no statistically significant differences of stroke/death when analyzing the general patient population.<sup>9, 14</sup> The second study also found no statistical difference in stroke/death/MI (OR, 1.4; 95% CI, 0.9–2.1; P=.18). Similarly, in three separate meta-analyses, no statistical differences were found between TCAR and CEA stroke or stroke/death outcomes in the overall population as well as the symptomatic and asymptomatic cohorts.<sup>13, 30, 31</sup> In terms of statistical significance, our findings likely reached significance due to the substantially larger pool of TCAR patients, which exceeded the largest meta-analysis study cohort (n=18,300) by nearly 32,000 patients. In a more recent and expansive VQI study involving 21,234 TCAR patients, CEA was associated with slightly lower odds of in-hospital stroke/death when compared with TCAR (1.7% vs 2.0%; aOR=0.82, 95% CI:0.72–0.95; P<.001), in line with our findings.<sup>7</sup> However, a significant limitation of these studies was the failure to delineate the preoperative symptom (amaurosis, hemispheric TIA, stroke) when adjusting for symptomatic status. Previous studies have demonstrated varying outcomes for patients presenting with amaurosis, hemispheric TIA, and stroke, with TCAR patients showing a higher incidence of preoperative strokes, while CEA patients tend to exhibit more preoperative ocular TIAs.<sup>32, 33, 34</sup> Consequently, amalgamating these diverse symptom statuses into a single ‘symptomatic’ variable could introduce bias into the analysis. Therefore, by preserving the distinction between these different symptom statuses during IPTW, we ensured a more nuanced and granular analysis.

With regards to in-hospital MI and CNI, we found that TCAR consistently had lower odds compared with CEA. In line with our findings, a systematic review and metanalysis comparing TCAR with CEA found that TCAR was associated with a lower incidence of MI and CNI.<sup>30</sup> While TCAR exhibited lower odds compared to CEA, it is noteworthy that the disparity in MI rates was relatively small. Additionally, although TCAR demonstrated a 2.0% lower CNI rate compared with CEA, the presence of a 0.3% CNI rate remains

significant, given the potential impact on patient quality of life.<sup>35</sup> As a result, our findings are consistent with previous studies, indicating that stroke/death outcomes, as well as MI and CNI are significant factors to consider when choosing between TCAR and CEA. However, further research is required to ascertain whether these statistically significant differences translate into clinically meaningful outcomes for patients, and more data on patient preferences are warranted.<sup>37, 38, 39, 40</sup> Additionally, physicians should account for anatomic considerations, procedure duration, anesthesia type, and procedural costs.<sup>24, 36</sup> With regards to anatomical considerations, CEA has been indicated for heavily calcified stenosis, angulated ICA, or low bifurcation where TCAR is not indicated. On the other hand, TCAR is often utilized for patients with a high bifurcation, prior endarterectomy, or other neck surgery/radiation that does not involve the base of the neck.<sup>41</sup>

When comparing TCAR with tfCAS and CEA with tfCAS, our study revealed worse outcomes for tfCAS, regardless of symptom status. Both arch manipulation and incomplete embolic protection are commonly cited factors thought to account for this increased risk of stroke/death associated with tfCAS.<sup>42</sup> While prior randomized control trials have found tfCAS stroke/death rates to be higher compared with TCAR and CEA, the low event rate and small patient numbers resulted in the inability to detect a statistically significant difference.<sup>19</sup> Nevertheless, this stroke/death difference may still be considered clinically important to patients and physicians. In a randomized control trial comparing tfCAS with CEA for patients with asymptomatic stenosis, the trial reported stenting was noninferior to endarterectomy, despite the higher stroke/death rates with tfCAS (2.9% vs 1.7%;  $P=.33$ ).<sup>44</sup> The Second Asymptomatic Carotid Surgery Trial (ACST-2) also found no statistical difference despite tfCAS having a stroke/death rate of 3.7% compared to 2.7% of CEA.<sup>45</sup>

Beyond considering the comparative stroke/death rates, it is also important to contextualize these outcomes over time and within the framework of acceptable standards. When investigating stroke/death rates over time, TCAR and CEA remain similar (with the exception of 2016 potentially attributable to the initiation of the TCAR Surveillance Project). In contrast, tfCAS stroke/death rates have remained high throughout the years with the exception of 2016 as well as a slight decrease in 2019. In terms of acceptable standards, the American Heart Association (AHA), Society for Vascular Surgery (SVS), and European guidelines recommend 30-day stroke/death rates to be <6% for symptomatic patients and <3% for asymptomatic patients based on the North American Symptomatic Carotid Endarterectomy Trial, European Carotid Surgery Trial, Asymptomatic Carotid Atherosclerosis Trial, and Asymptomatic Carotid Surgery Trial.<sup>46, 47, 48, 49, 45</sup> However, prior studies have shown that at least 1/3 of 30-day stroke/death events occur post-discharge, and therefore, these in-hospital rates are underestimating adverse outcomes.<sup>50, 51</sup> Consequently, there is a growing consensus that acceptable in-hospital stroke/death rates should be reevaluated, with suggestions proposing ~4% for symptomatic cases and ~2% for asymptomatic cases.<sup>52, 53, 54, 55, 56</sup> In this context, our results indicate that while both CEA and TCAR fall within the threshold of acceptability, tfCAS would not meet these benchmarks for either symptomatic or asymptomatic patients. These findings substantiate existing safety concerns surrounding tfCAS and emphasize the importance of careful patient selection.

While tfCAS did have the lowest rates of CNI compared to both TCAR and CEA, the relatively larger stroke/death rates support the prioritization of TCAR and CEA, with the use of tfCAS for select indications such as patients with a tracheostomy, prior neck surgery, significant plaque at the TCAR sheath access site, or contralateral CNI.<sup>57, 58</sup> Nevertheless, CMS has expanded tfCAS indications to include asymptomatic patients and patients at standard risk for adverse events from CEA without a need for registry participation or center certification. The label expansion will also allow a larger number of patients and physicians access to tfCAS.<sup>59, 60</sup> The current analysis highlights the importance of continued investigation with registry data collection, randomized clinical trials such as CREST 2, and observational studies to ensure patient safety.<sup>61</sup>

While our study brings additional insight to the field through analysis of a larger TCAR population on the comparative safety of TCAR, tfCAS, and CEA, it is important to acknowledge the following limitations. First, our study was observational and subjected to the limitations inherent in a retrospective study, including the presence of unmeasured confounding. Furthermore, the lack of data concerning the nature of carotid lesions, stroke severity, Instructions for Use status, and the clinical significance of CNI, MI, and stroke on quality of life precludes further analysis on these topics. Additionally, our findings predominantly address in-hospital outcomes, limiting our ability to draw conclusions regarding long-term outcomes and recurrence rates. VQI captures >95% of TCAR patients, however, many CEA and tfCAS patients are not included in the VQI dataset, potentially limiting the comprehensiveness of the results in reflecting national practice patterns and variations across different specialties. Moreover, physician information is also blinded, which prevents an analysis of outcomes by specialty (e.g., vascular surgery vs interventional radiology vs cardiology). While specific breakdowns are not provided, the VQI does include a substantial number of cardiologists and interventional radiologists, particularly since the merger of the ACC NCDR dataset into the VQI in 2021. Furthermore, during the initial years of our analysis, TCAR and tfCAS were exclusively recommended for high-risk patients. Consequently, these early years may have led to a higher occurrence of adverse outcomes compared to CEA. To control for this, we incorporated the year of surgery into our model. Of note, this potential bias would not hamper a comparison of TCAR with tfCAS. Nevertheless, the recent increase in TCAR utilization and respective decrease in tfCAS use may impact outcomes, further underscoring the importance of continuously updating clinical data to reflect current practice patterns. Additionally, it is noteworthy that not all physicians may possess expertise in all three techniques, and some patients may be ineligible for a certain procedure. Consequently, continuous research is needed to address the current limitations of the VQI to facilitate the selection of the most appropriate revascularization technique tailored to individual clinical or regional contexts. A randomized trial comparing CEA, TCAR, and tfCAS, coupled with ongoing monitoring of outcomes using registry data, holds significance in optimizing patient care. Nonetheless, the present findings serve to guide shared decision-making discussions when exploring treatment options.

## Conclusion

In the VQI registry, TCAR had clinically similar rates of stroke/death compared to CEA. On the other hand, tfCAS in-hospital outcomes proved inferior to the other two procedures, with

higher rates of stroke/death. As a result, this 7-year analysis underscores the comparability of TCAR and CEA, highlights the relatively poor outcomes of tFCAS, and serves as a valuable resource for shared decision-making in revascularization.

## Grant support

Research reported in this publication was supported by *NIH offices* of the National Institutes of Health under award numbers **5T35HL110843-10**, **5T32HL007734-29**, and **1UM1TR004408-01**.

## References

1. Cole TS, Mezher AW, Catapano JS, Godzik J, Baranoski JF, Nakaji P, et al. Nationwide Trends in Carotid Endarterectomy and Carotid Artery Stenting in the Post-CREST Era. *Stroke*. 2020;51(2):579–587. doi:10.1161/STROKEAHA.119.027388 [PubMed: 31847750]
2. Saleem T, Baril DT. Carotid Artery Stenting. In: StatPearls. StatPearls Publishing; 2023. Accessed August 9, 2023. <http://www.ncbi.nlm.nih.gov/books/NBK470541/>
3. Schermerhorn ML, Liang P, Eldrup-Jorgensen J, Cronenwett JL, Nolan BW, Kashyap VS, et al. Association of Transcarotid Artery Revascularization vs Transfemoral Carotid Artery Stenting With Stroke or Death Among Patients With Carotid Artery Stenosis. *JAMA*. 2019;322(23):2313–2322. doi:10.1001/jama.2019.18441 [PubMed: 31846015]
4. Premarket Approval (PMA). Accessed August 9, 2023. <https://www.accessdata.fda.gov/scripts/cdrh/cfdocs/cfpma/pma.cfm?id=P140026S016>
5. Kwolek CJ, Jaff MR, Leal JI, Hopkins LN, Shah RM, Hanover TM, et al. Results of the ROADSTER multicenter trial of transcarotid stenting with dynamic flow reversal. *J Vasc Surg*. 2015;62(5):1227–1234. doi:10.1016/j.jvs.2015.04.460 [PubMed: 26506270]
6. Malas M, Nejm BJ, Kwolek CJ, Leal Lorenzo JI, Hanover T, Mehta M, et al. One-Year Results of the ROADSTER Multicenter Trial of Transcarotid Stenting With Dynamic Flow Reversal. *Journal of Vascular Surgery*. 2018;67(1):e5. doi:10.1016/j.jvs.2017.10.012
7. Columbo JA, Martinez-Cambor P, Stone DH, Goodney PP, O'Malley AJ. Procedural Safety Comparison Between Transcarotid Artery Revascularization, Carotid Endarterectomy, and Carotid Stenting: Perioperative and 1-Year Rates of Stroke or Death. *J Am Heart Assoc*. 2022;11(19):e024964. doi:10.1161/JAHA.121.024964 [PubMed: 36172943]
8. Kibrik P, Stonko DP, Alsheekh A, Holscher C, Zarkowsky D, Abularrage CJ, et al. Association of Carotid Revascularization Approach with Perioperative Outcomes Based on Symptom Status and Degree of Stenosis among Octogenarians R3WC: 341/3181. *J Vasc Surg*. 2022;76(3):769–777.e2. doi:10.1016/j.jvs.2022.04.027 [PubMed: 35643202]
9. Malas MB, Dakour-Aridi H, Kashyap VS, Eldrup-Jorgensen J, Wang GJ, Motaganahalli RL, et al. TransCarotid Revascularization With Dynamic Flow Reversal Versus Carotid Endarterectomy in the Vascular Quality Initiative Surveillance Project. *Ann Surg*. 2022;276(2):398–403. doi:10.1097/SLA.0000000000004496 [PubMed: 32941280]
10. Mehta A, Patel P, Bajakian D, Schutzer R, Morrissey N, Garg K, et al. Transcarotid Artery Revascularization Versus Carotid Endarterectomy and Transfemoral Stenting in Octogenarians. *Journal of Vascular Surgery*. 2020;72(3):e284. doi:10.1016/j.jvs.2020.06.030
11. Silk Road Medical's Enroute TCAR System Approved for Expanded Indication of Standard-Risk Patients - Endovascular Today. Accessed August 19, 2023. <https://evtoday.com/news/silk-road-medicals-enroute-tcar-system-approved-for-expanded-indication-of-standard-risk-patients>
12. Stonko DP, Goldsborough E III, Kibrik P, Zhang G, Holscher CM, Hicks CW. Use of Transcarotid Artery Revascularization, Transfemoral Carotid Artery Stenting, and Carotid Endarterectomy in the US From 2015 to 2019. *JAMA Network Open*. 2022;5(9):e2231944. doi:10.1001/jamanetworkopen.2022.31944 [PubMed: 36112371]
13. Naazie IN, Cui CL, Osaghae I, Murad MH, Schermerhorn M, Malas MB. A Systematic Review and Meta-Analysis of Transcarotid Artery Revascularization with Dynamic Flow Reversal Versus Transfemoral Carotid Artery Stenting and Carotid Endarterectomy. *Ann Vasc Surg*. 2020;69:426–436. doi:10.1016/j.avsg.2020.05.070 [PubMed: 32505684]

14. Schermerhorn ML, Liang P, Dakour-Aridi H, Kashyap VS, Wang GJ, Nolan BW, et al. In-hospital outcomes of transcarotid artery revascularization and carotid endarterectomy in the Society for Vascular Surgery Vascular Quality Initiative. *Journal of Vascular Surgery*. 2020;71(1):87–95. doi:10.1016/j.jvs.2018.11.029 [PubMed: 31227410]
15. Liang P, Cronenwett JL, Secemsky EA, Eldrup-Jorgensen J, Malas MB, Wang GJ, et al. Risk of Stroke, Death, and Myocardial Infarction Following Transcarotid Artery Revascularization vs Carotid Endarterectomy in Patients With Standard Surgical Risk. *JAMA Neurol*. 2023;80(5):437–444. doi:10.1001/jamaneurol.2023.0285 [PubMed: 36939697]
16. Cui CL, Dakour-Aridi H, Lu JJ, Yei KS, Schermerhorn ML, Malas MB. In-Hospital Outcomes of Urgent, Early, or Late Revascularization for Symptomatic Carotid Artery Stenosis. *Stroke*. 2022;53(1):100–107. doi:10.1161/STROKEAHA.120.032410 [PubMed: 34872337]
17. Yei KS, Janssen C, Elsayed N, Naazie I, Sedrakyan A, Malas MB. Long-term outcomes of carotid endarterectomy vs transfemoral carotid stenting in a Medicare-matched database. *J Vasc Surg*. Published online August 25, 2023:S0741–5214(23)01932–8. doi:10.1016/j.jvs.2023.08.118
18. Galyfos G, Sachsamanis G, Anastasiadou C, Sachmpazidis I, Kikiras K, Kastrisios G, et al. Carotid Endarterectomy versus Carotid Stenting or Best Medical Treatment in Asymptomatic Patients with Significant Carotid Stenosis: A meta-analysis. *Cardiovascular Revascularization Medicine*. 2019;20(5):413–423. doi:10.1016/j.carrev.2018.07.003 [PubMed: 30057288]
19. Müller MD, Lyrer PA, Brown MM, Bonati LH. Carotid Artery Stenting Versus Endarterectomy for Treatment of Carotid Artery Stenosis. *Stroke*. 2021;52(1):e3–e5. doi:10.1161/STROKEAHA.120.030521 [PubMed: 33370200]
20. Mantese VA, Timaran CH, Chiu D, Begg RJ, Brott TG. The Carotid Revascularization Endarterectomy versus Stenting Trial (CREST) - Stenting versus Carotid Endarterectomy for Carotid Disease. *Stroke*. 2010;41(10 Suppl):S31–S34. doi:10.1161/STROKEAHA.110.595330 [PubMed: 20876500]
21. Delgado C, Baweja M, Burrows NR, Crews DC, Eneanya ND, Gadegbeku CA, et al. Reassessing the Inclusion of Race in Diagnosing Kidney Diseases: An Interim Report From the NKF-ASN Task Force. *Am J Kidney Dis*. 2021;78(1):103–115. doi:10.1053/j.ajkd.2021.03.008 [PubMed: 33845065]
22. Dakour-Aridi H, Giuliano K, Locham S, Dang T, Siracuse JJ, Malas MB. Perioperative blood transfusion in anemic patients undergoing elective endovascular abdominal aneurysm repair. *Journal of Vascular Surgery*. 2020;71(1):75–85. doi:10.1016/j.jvs.2019.02.065 [PubMed: 31176635]
23. Timaran CH, McKinsey JF, Schneider PA, Littooy F. Reporting standards for carotid interventions from the Society for Vascular Surgery. *J Vasc Surg*. 2011;53(6):1679–1695. doi:10.1016/j.jvs.2010.11.122 [PubMed: 21609800]
24. Glaser J, Kuwayama D, Stone D, Schanzer A, Eldrup-Jorgensen J, Powell R, et al. Factors that determine the length of stay after carotid endarterectomy represent opportunities to avoid financial losses. *J Vasc Surg*. 2014;60(4):966–972.e1. doi:10.1016/j.jvs.2014.03.292 [PubMed: 24865784]
25. Fourth Universal Definition of Myocardial Infarction. American College of Cardiology. Accessed April 26, 2024. <https://www.acc.org/Latest-in-Cardiology/ten-points-to-remember/2019/01/21/14/44/http%3a%2f%2fwww.acc.org%2fLatest-in-Cardiology%2ften-points-to-remember%2f2019%2f01%2f21%2f14%2f44%2fFourth-Universal-Definition-of-Myocardial-Infarction>
26. Stuart EA, Huskamp HA, Duckworth K, Simmons J, Song Z, Chernew M, et al. Using propensity scores in difference-in-differences models to estimate the effects of a policy change. *Health Serv Outcomes Res Methodol*. 2014;14(4):166–182. doi:10.1007/s10742-014-0123-z [PubMed: 25530705]
27. Griffin BA, Ramchand R, Almirall D, Slaughter ME, Burgette LF, McCaffery DF. Estimating the causal effects of cumulative treatment episodes for adolescents using marginal structural models and inverse probability of treatment weighting. *Drug Alcohol Depend*. 2014;136:69–78. doi:10.1016/j.drugalcdep.2013.12.017 [PubMed: 24440050]
28. Buck DB, Soden PA, Deery SE, Zettervall SL, Ultee KHJ, Landon BE, et al. Comparison of Endovascular Stent Grafts for Abdominal Aortic Aneurysm Repair in Medicare Beneficiaries. *Ann Vasc Surg*. 2018;47:31–42. doi:10.1016/j.avsg.2017.08.021 [PubMed: 28890065]

29. Wu WW, Liang P, O'Donnell TFX, Swerdlow NJ, Li C, Wyers MC, et al. Anatomic Eligibility for Transcarotid Artery Revascularization and Transfemoral Carotid Artery Stenting. *J Vasc Surg.* 2019;69(5):1452–1460. doi:10.1016/j.jvs.2018.11.051 [PubMed: 30853384]
30. Gao J, Chen Z, Kou L, Zhang H, Yang Y. The Efficacy of Transcarotid Artery Revascularization With Flow Reversal System Compared to Carotid Endarterectomy: A Systematic Review and Meta-Analysis. *Front Cardiovasc Med.* 2021;8:695295. doi:10.3389/fcvm.2021.695295 [PubMed: 34869622]
31. Wu H, Wang Z, Li M, Sun P, Wei S, Xie B, et al. Outcomes of transcarotid artery revascularization: A systematic review. *Interv Neuroradiol.* Published online August 29, 2022:15910199221123283. doi:10.1177/15910199221123283
32. Solomon Y, Rastogi V, Marcaccio CL, Patel PB, Wang GJ, Malas MB, et al. Outcomes after transcarotid artery revascularization stratified by preprocedural symptom status. *J Vasc Surg.* 2022;76(5):1307–1315.e1. doi:10.1016/j.jvs.2022.05.024 [PubMed: 35798281]
33. Solomon Y, Varkevisser RRB, Swerdlow NJ, Li C, Liang P, Siracuse JJ, et al. Outcomes after transfemoral carotid artery stenting stratified by preprocedural symptom status. *J Vasc Surg.* 2021;73(6):2021–2029. doi:10.1016/j.jvs.2020.11.031 [PubMed: 33278538]
34. Pothof AB, Zwanenburg ES, Deery SE, O'Donnell TFX, de Borst GJ, Schermerhorn ML. An update on the incidence of perioperative outcomes after carotid endarterectomy, stratified by type of preprocedural neurologic symptom. *J Vasc Surg.* 2018;67(3):785–792. doi:10.1016/j.jvs.2017.07.132 [PubMed: 29074118]
35. Hye RJ, Mackey A, Hill MD, Voeks J, Cohen DJ, Wang K, et al. Incidence, Outcomes and Effect on Quality of Life of Cranial Nerve Injury (CNI) in the Carotid Revascularization Endarterectomy versus Stenting Trial (CREST). *J Vasc Surg.* 2015;61(5):1208–1215. doi:10.1016/j.jvs.2014.12.039 [PubMed: 25770984]
36. Kanitra JJ, Graham IA, Hayward RD, Granger DK, Berg RA, Haouilou JC. Estimated Cost of Transcarotid Arterial Revascularization Compared With Carotid Endarterectomy and Transfemoral Carotid Stenting. *Cureus.* 14(3):e23539. doi:10.7759/cureus.23539
37. Columbo JA, Martinez-Camblor P, O'Malley AJ, Stone DH, Kashyap VS, Powell RJ, et al. Association of Adoption of Transcarotid Artery Revascularization With Center-Level Perioperative Outcomes. *JAMA Netw Open.* 2021;4(2):e2037885. doi:10.1001/jamanetworkopen.2020.37885 [PubMed: 33616666]
38. Lal BK, Beach KW, Roubin GS, Lutsep HL, Moore WS, Malas MB, et al. Restenosis after carotid artery stenting and endarterectomy: a secondary analysis of CREST, a randomised controlled trial. *Lancet Neurol.* 2012;11(9):755–763. doi:10.1016/S1474-4422(12)70159-X [PubMed: 22857850]
39. Howard DPJ, Gaziano L, Rothwell PM. Risk of stroke in relation to degree of asymptomatic carotid stenosis: a population-based cohort study, systematic review, and meta-analysis. *The Lancet Neurology.* 2021;20(3):193–202. doi:10.1016/S1474-4422(20)30484-1 [PubMed: 33609477]
40. Turan TN, Voeks JH, Chimowitz MI, Roldan A, LeMatty T, Haley W, et al. Rationale, Design, and Implementation of Intensive Risk Factor Treatment in the CREST2 Trial. *Stroke.* 2020;51(10):2960–2971. doi:10.1161/STROKEAHA.120.030730 [PubMed: 32951538]
41. Fan W, Shi W, Lu S, Guo W, Tong J, Tan J, et al. Analysis of the anatomic eligibility for transcarotid artery revascularization in Chinese patients who underwent carotid endarterectomy and transfemoral carotid artery stenting. *Front Cardiovasc Med.* 2023;9:1045598. doi:10.3389/fcvm.2022.1045598 [PubMed: 36684575]
42. Naazie IN, Cui CL, Osaghae I, Murad MH, Schermerhorn M, Malas MB. A Systematic Review and Meta-Analysis of Transcarotid Artery Revascularization with Dynamic Flow Reversal Versus Transfemoral Carotid Artery Stenting and Carotid Endarterectomy. *Annals of Vascular Surgery.* 2020;69:426–436. doi:10.1016/j.avsg.2020.05.070 [PubMed: 32505684]
43. Cui CL, Zarrintan S, Marmor RA, Nichols J, Cajas-Monson L, Malas M. Performance of Carotid Revascularization Procedures as Modified by Sex. *Annals of Vascular Surgery.* 2022;81:171–182. doi:10.1016/j.avsg.2021.08.051 [PubMed: 34752853]
44. Rosenfield K, Matsumura JS, Chaturvedi S, Riles T, Ansel GM, Metzger DC, et al. Randomized Trial of Stent versus Surgery for Asymptomatic Carotid Stenosis. *N Engl J Med.* 2016;374(11):1011–1020. doi:10.1056/NEJMoa1515706 [PubMed: 26886419]

45. Halliday A, Bulbulia R, Bonati LH, Chester J, Craddock-Bamford A, Peto R, et al. Second asymptomatic carotid surgery trial (ACST-2): a randomised comparison of carotid artery stenting versus carotid endarterectomy. *The Lancet*. 2021;398(10305):1065–1073. doi:10.1016/S0140-6736(21)01910-3
46. Randomised trial of endarterectomy for recently symptomatic carotid stenosis: final results of the MRC European Carotid Surgery Trial (ECST). *The Lancet*. 1998;351(9113):1379–1387. doi:10.1016/S0140-6736(97)09292-1
47. Beneficial Effect of Carotid Endarterectomy in Symptomatic Patients with High-Grade Carotid Stenosis. *New England Journal of Medicine*. 1991;325(7):445–453. doi:10.1056/NEJM199108153250701 [PubMed: 1852179]
48. Walker MD, Marler JR, Goldstein M, Grady PA, Toole JF, Baker WH, et al. Endarterectomy for Asymptomatic Carotid Artery Stenosis. *JAMA*. 1995;273(18):1421–1428. doi:10.1001/jama.1995.03520420037035 [PubMed: 7723155]
49. Moore WS, Barnett H j. m., Beebe HG, Bernstein EF, Brener BJ, Brott T, et al. Guidelines for Carotid Endarterectomy. *Stroke*. 1995;26(1):188–201. doi:10.1161/01.STR.26.1.188 [PubMed: 7839390]
50. Liang P, Solomon Y, Swerdlow NJ, Li C, Varkevisser RRB, de Guerre LEVM, et al. In-hospital Outcomes Alone Underestimate Rates of 30-Day Major Adverse Events following Carotid Artery Stenting. *J Vasc Surg*. 2020;71(4):1233–1241. doi:10.1016/j.jvs.2019.06.201 [PubMed: 32063441]
51. Fokkema M, Bensley R, Lo R, Hamdan A, Wyers M, Moll F, et al. In-hospital versus post discharge adverse events following carotid endarterectomy. *J Vasc Surg*. 2013;57(6):1568–1575.e3. doi:10.1016/j.jvs.2012.11.072 [PubMed: 23388394]
52. Schneider PA, Naylor AR. Asymptomatic carotid artery stenosis—Medical therapy alone versus medical therapy plus carotid endarterectomy or stenting. *Journal of Vascular Surgery*. 2010;52(2):499–507. doi:10.1016/j.jvs.2010.05.063 [PubMed: 20670781]
53. Naylor R, Rantner B, Ancetti S, Borst GJ de, Carlo MD, Halliday A, et al. Editor’s Choice – European Society for Vascular Surgery (ESVS) 2023 Clinical Practice Guidelines on the Management of Atherosclerotic Carotid and Vertebral Artery Disease. *European Journal of Vascular and Endovascular Surgery*. 2023;65(1):7–111. doi:10.1016/j.ejvs.2022.04.011 [PubMed: 35598721]
54. Bonati LH, Kakkos S, Berkefeld J, de Borst GJ, Bulbulia R, Halliday A, et al. European Stroke Organisation guideline on endarterectomy and stenting for carotid artery stenosis. *European Stroke Journal*. 2021;6(2):I–XLVII. doi:10.1177/23969873211012121
55. Gray WA, Chaturvedi S, Verta P. Thirty-Day Outcomes for Carotid Artery Stenting in 6320 Patients From 2 Prospective, Multicenter, High-Surgical-Risk Registries. *Circulation: Cardiovascular Interventions*. 2009;2(3):159–166. doi:10.1161/CIRCINTERVENTIONS.108.823013 [PubMed: 20031712]
56. Geiger JT, Fleming F, Iannuzzi JC, Stoner M, Doyle A. Guideline Compliant Minimum Asymptomatic Carotid Endarterectomy Surgeon and Hospital Volume Cutoffs. *Annals of Vascular Surgery*. 2023;97:129–138. doi:10.1016/j.avsg.2023.07.089 [PubMed: 37454899]
57. Kumins NH, King AH, Ambani RN, Thomas JP, Kim AH, Augustin G, et al. Anatomic criteria in the selection of treatment modality for atherosclerotic carotid artery disease. *J Vasc Surg*. 2020;72(4):1395–1404. doi:10.1016/j.jvs.2020.01.041 [PubMed: 32145991]
58. Vitek JJ, Roubin GS, Al-Mubarek N, New G, Iyer SS. Carotid Artery Stenting: Technical Considerations. *AJNR Am J Neuroradiol*. 2000;21(9):1736–1743. [PubMed: 11039358]
59. NCA - Carotid Artery Stenting (CAG-00085R) - Decision Memo. Accessed August 2, 2023. <https://www.cms.gov/medicare-coverage-database/view/ncacal-decision-memo.aspx?proposed=N&NCAId=157>
60. Silk Road Medical Announces FDA Approval of Expanded Indications for the ENROUTE<sup>®</sup> Transcarotid Stent System - Silk Road Medical, Inc. Accessed August 2, 2023. <https://investors.silkroadmed.com/news-releases/news-release-details/silk-road-medical-announces-fda-approval-expanded-indications/>

61. Mott M, Koroshetz W, Wright CB. CREST-2: Identifying the Best Method of Stroke Prevention for Carotid Artery Stenosis. *Stroke*. 2017;48(5):e130–e131. doi:10.1161/STROKEAHA.117.016051 [PubMed: 28386040]

Author Manuscript

Author Manuscript

Author Manuscript

Author Manuscript

**ARTICLE HIGHLIGHTS****Type of Research:**


Retrospective analysis of prospectively collected data from the Vascular Quality Initiative

**Key Findings:**

From a database encompassing over 50,000 transcrotid artery revascularization, 25,000 transfemoral carotid artery stenting, and 120,000 carotid endarterectomy patients, our analysis revealed transfemoral carotid artery stenting was associated with higher rates of in-hospital stroke/death compared with both transcrotid artery revascularization and carotid endarterectomy. Given the relatively smaller differences in stroke/death outcomes between transcrotid artery revascularization and endarterectomy, determining clinical significance becomes paramount in assessing the overall risk-benefit profile of these two interventions.

**Take home Message:**

While transfemoral carotid artery stenting may be beneficial for select patients, the decision between carotid endarterectomy and transfemoral carotid artery revascularization should be multifactorial.



**Figure 1:**  
Adjusted rates of stroke/death overtime for TCAR, tfCAS, and CEA  
\*Note: 2023 data includes up until mid-August

**Table I:**

Baseline characteristics of patients undergoing CEA, TCAR, or tfCAS for carotid stenosis

	Baseline comparison				Adjusted Rates			
	TCAR	TFCAS	CEA	<i>P value</i>	TCAR	TFCAS	CEA	<i>P value</i>
N	50068	25361	122737					
Age mean (SD) <sup>1</sup>	73.5 (8.59)	70.5 (9.45)	71.0 (8.90)	< .001	72.2 (0.05)	72.8 (0.06)	71.1 (0.03)	.774
Female Sex	37.3%	36.2%	39.4%	< .001	37.3%	36.2%	39.4%	<.001
White Race	89.9%	88.7%	89.8%	< .001	89.5%	89.3%	89.9%	<.001
Hispanic Ethnicity	4.4%	4.0%	3.3%	< .001	4.5%	4.1%	3.3%	<.001
Symptomatic	24.7%	36.9%	29.6%	< .001	27.6%	32.1%	29.3%	<.001
Diabetes	39.0%	39.5%	37.1%	< .001	39.1%	39.7%	37.1%	<.001
CAD <sup>2</sup>	53.7%	50.8%	39.6%	< .001	52.9%	53.8%	39.7%	<.001
CHF <sup>3</sup>	16.7%	16.8%	12.1%	< .001	16.4%	18.1%	12.2%	<.001
CKD <sup>4</sup>	38.8%	34.9%	24.0%	< .001	37.8%	37.7%	24.7%	<.001
COPD <sup>5</sup>	24.7%	25.3%	22.9%	< .001	25.0%	24.9%	22.9%	<.001
CABG/PCI <sup>6</sup>	38.6%	36.9%	33.4%	< .001	37.3%	39.8%	33.6%	<.001
Prior Contralateral CEA/CAS	16.2%	15.1%	12.4%	< .001	15.4%	16.7%	12.5%	<.001
Prior Ipsilateral CEA/CAS	12.6%	17.2%	1.4%	< .001	12.4%	17.4%	1.5%	<.001
Betablocker	53.7%	50.8%	49.8%	< .001	53.3%	52.1%	49.9%	<.001
ACE-Inhibitor	54.2%	49.8%	53.6%	< .001	53.8%	50.3%	53.7%	<.001
Aspirin	89.8%	84.5%	83.9%	< .001	89.9%	84.0%	83.9%	<.001
Anticoagulant	4.0%	4.5%	12.8%	< .001	3.9%	4.3%	12.9%	<.001
Antiplatelets	89.1%	76.8%	38.5%	< .001	88.8%	76.8%	38.6%	<.001
Statin	90.2%	83.2%	85.9%	< .001	90.2%	83.6%	85.9%	<.001
Ipsilateral Stenosis >80%	47.0%	52.1%	44.6%	< .001	48.4%	52.2%	45.6%	<.001

<sup>1</sup> Standard Deviation (SD)<sup>2</sup> Coronary Artery Disease (CAD)<sup>3</sup> Congestive Heart Failure (CHF)<sup>4</sup> Chronic Kidney Disease (CKD)<sup>5</sup> Chronic Obstructive Pulmonary Disorder (COPD)<sup>6</sup> Coronary Artery Bypass Grafting / Percutaneous Coronary Intervention (CABG/PCI)

**Table II:**

Postoperative outcomes for all carotid stenosis patients

				TFCAS vs TCAR		TFCAS vs CEA		CEA vs TCAR	
	TCAR	TFCAS	CEA	aOR [CI]	P value	aOR [CI]	P value	aOR [CI]	P value
In-hospital Stroke	1.3%	2.1%	1.1%	1.58 (1.39–1.79)	<.001	1.88 (1.68–2.10)	<.001	0.84 (0.76–0.93)	<.001
In-hospital Death	0.4%	1.1%	0.3%	2.80 (2.29–3.42)	<.001	3.66 (3.10–4.32)	<.001	0.76 (0.64–0.92)	.004
Perioperative MI	0.5%	0.4%	0.6%	0.88 (0.69–1.13)	.3	0.67 (0.54–0.84)	<.001	1.31 (1.13–1.54)	<.001
Stroke or Death	1.6%	2.9%	1.3%	1.84 (1.65–2.06)	<.001	2.21 (2.01–2.43)	<.001	0.83 (0.76–0.91)	<.001
Stroke, Death, MI	2.0%	3.2%	1.9%	1.62 (1.46–1.80)	<.001	1.72 (1.58–1.88)	<.001	0.94 (0.87–1.02)	.14
Length of Stay > 2 days	29.6%	35.5%	30.1%	1.31 (1.27–1.36)	<.001	1.28 (1.24–1.32)	<.001	1.02 (1.00–1.05)	.047
Cranial Nerve Injury	0.3%	0.0%	2.3%	0.00 (0.00–0.00)	<.001	0.00 (0.00–0.00)	<.001	9.42 (7.78–11.4)	<.001
Bleeding	0.8%	0.4%	1.0%	0.52 (0.41–0.67)	<.001	0.38 (0.30–0.47)	<.001	1.38 (1.22–1.57)	<.001

\* IPTW accounted for the following variables: age, sex, race, BMI, hypertension, diabetes, CAD, CHF, preoperative smoking, COPD, renal dysfunction, anemia, symptomatic amaurosis, symptomatic hemispheric TIA, symptomatic stroke, CABG / PCI, prior contralateral CEA/CAS, % ipsilateral and contralateral occlusion, and surgery year.

**Table III:**

Postoperative outcomes for symptomatic carotid stenosis patients

				TFCAS vs TCAR		TFCAS vs CEA		CEA vs TCAR	
	TCAR	TFCAS	CEA	aOR [CI]	P value	aOR [CI]	P value	aOR [CI]	P value
In-hospital Stroke	2.1%	3.0%	1.8%	1.46 (1.21–1.76)	<.001	1.73 (1.48–2.02)	<.001	0.84 (0.72–0.98)	.029
In-hospital Death	0.6%	1.9%	0.5%	3.35 (2.50–4.48)	<.001	4.06 (3.26–5.05)	<.001	0.83 (0.62–1.10)	.2
Perioperative MI	0.4%	0.6%	0.6%	1.39 (0.93–2.07)	.10	0.97 (0.71–1.32)	.8	1.44 (1.05–1.96)	.022
Stroke or Death	2.4%	4.4%	2.1%	1.87 (1.59–2.21)	<.001	2.18 (1.91–2.48)	<.001	0.86 (0.75–0.99)	.039
Stroke, Death, MI	2.8%	4.8%	2.6%	1.78 (1.53–2.08)	<.001	1.89 (1.67–2.14)	<.001	0.94 (0.83–1.08)	.4
Length of Stay > 2 days	41.1%	52.2%	42.9%	1.56 (1.47–1.66)	<.001	1.45 (1.38–1.53)	<.001	1.08 (1.03–1.12)	.001
Cranial Nerve Injury	0.3%	0.0%	2.8%	0.00 (0.00–0.00)	<.001	0.00 (0.00–0.00)	<.001	8.60 (6.20–11.9)	<.001
Bleeding	0.8%	0.5%	1.3%	0.58 (0.39–0.87)	.009	0.36 (0.25–0.51)	<.001	1.61 (1.28–2.04)	<.001

**Table IV:**

Postoperative outcomes for asymptomatic carotid stenosis patients

				TFCAS vs TCAR		TFCAS vs CEA		CEA vs TCAR	
	TCAR	TFCAS	CEA	aOR [CI]	P value	aOR [CI]	P value	aOR [CI]	P value
In-hospital Stroke	1.1%	1.7%	0.9%	1.57 (1.32–1.86)	<.001	1.94 (1.66–2.27)	<.001	0.81 (0.71–0.91)	<.001
In-hospital Death	0.3%	0.7%	0.2%	2.13 (1.59–2.85)	<.001	3.07 (2.37–3.98)	<.001	0.69 (0.54–0.88)	.003
Perioperative MI	0.5%	0.3%	0.6%	0.67 (0.48–0.94)	.020	0.53 (0.39–0.73)	<.001	1.26 (1.06–1.51)	.011
Stroke or Death	1.3%	2.1%	1.0%	1.68 (1.44–1.96)	<.001	2.15 (1.87–2.47)	<.001	0.78 (0.70–0.88)	<.001
Stroke, Death, MI	1.7%	2.4%	1.5%	1.42 (1.23–1.64)	<.001	1.56 (1.37–1.77)	<.001	0.91 (0.82–1.01)	.067
Length of Stay > 2 days	25.3%	27.6%	24.7%	1.12 (1.07–1.18)	<.001	1.16 (1.11–1.21)	<.001	0.97 (0.94–1.00)	.027
Cranial Nerve Injury	0.2%	0.0%	2.2%	0.00 (0.00–0.00)	<.001	0.00 (0.00–0.00)	<.001	9.67 (7.64–12.3)	<.001
Bleeding	0.7%	0.4%	0.9%	0.49 (0.36–0.68)	<.001	0.38 (0.29–0.51)	<.001	1.29 (1.12–1.50)	<.001