

## ORIGINAL ARTICLE

# Transcatheter Valve Repair in Heart Failure with Moderate to Severe Mitral Regurgitation

S.D. Anker, T. Friede, R.-S. von Bardeleben, J. Butler, M.-S. Khan, M. Diek, J. Heinrich, M. Geyer, M. Placzek, R. Ferrari, W.T. Abraham, O. Alfieri, A. Auricchio, A. Bayes-Genis, J.G.F. Cleland, G. Filippatos, F. Gustafsson, W. Haverkamp, M. Kelm, K.-H. Kuck, U. Landmesser, A.P. Maggioni, M. Metra, V. Ninios, M.C. Petrie, T. Rassaf, F. Ruschitzka, U. Schäfer, P.C. Schulze, K. Spargias, A. Vahanian, J.L. Zamorano, A. Zeiher, M. Karakas, F. Koehler, M. Lainscak, A. Öner, N. Mezilis, E.K. Theofilogiannakos, I. Ninios, M. Chrissoheris, P. Kourkouveli, K. Papadopoulos, G. Smolka, W. Wojakowski, K. Reczuch, F.J. Pinto, Ł. Wiewiórka, Z. Kalarus, M. Adamo, E. Santiago-Vacas, T.F. Ruf, M. Gross, J. Tongers, G. Hasenfuss, W. Schillinger, and P. Ponikowski, for the RESHAPE-HF2 Investigators\*

## ABSTRACT

**BACKGROUND**

Whether transcatheter mitral-valve repair improves outcomes in patients with heart failure and functional mitral regurgitation is uncertain.

**METHODS**

We conducted a randomized, controlled trial involving patients with heart failure and moderate to severe functional mitral regurgitation from 30 sites in nine countries. The patients were assigned in a 1:1 ratio to either transcatheter mitral-valve repair and guideline-recommended medical therapy (device group) or medical therapy alone (control group). The three primary end points were the rate of the composite of first or recurrent hospitalization for heart failure or cardiovascular death during 24 months; the rate of first or recurrent hospitalization for heart failure during 24 months; and the change from baseline to 12 months in the score on the Kansas City Cardiomyopathy Questionnaire—Overall Summary (KCCQ-OS; scores range from 0 to 100, with higher scores indicating better health status).

**RESULTS**

A total of 505 patients underwent randomization: 250 were assigned to the device group and 255 to the control group. At 24 months, the rate of first or recurrent hospitalization for heart failure or cardiovascular death was 37.0 events per 100 patient-years in the device group and 58.9 events per 100 patient-years in the control group (rate ratio, 0.64; 95% confidence interval [CI], 0.48 to 0.85;  $P=0.002$ ). The rate of first or recurrent hospitalization for heart failure was 26.9 events per 100 patient-years in the device group and 46.6 events per 100 patient-years in the control group (rate ratio, 0.59; 95% CI, 0.42 to 0.82;  $P=0.002$ ). The KCCQ-OS score increased by a mean ( $\pm$ SD) of  $21.6\pm 26.9$  points in the device group and  $8.0\pm 24.5$  points in the control group (mean difference, 10.9 points; 95% CI, 6.8 to 15.0;  $P<0.001$ ). Device-specific safety events occurred in 4 patients (1.6%).

**CONCLUSIONS**

Among patients with heart failure with moderate to severe functional mitral regurgitation who received medical therapy, the addition of transcatheter mitral-valve repair led to a lower rate of first or recurrent hospitalization for heart failure or cardiovascular death and a lower rate of first or recurrent hospitalization for heart failure at 24 months and better health status at 12 months than medical therapy alone. (Funded by Abbott Laboratories; RESHAPE-HF2 ClinicalTrials.gov number, NCT02444338.)

The authors' full names, academic degrees, and affiliations are listed in the Appendix. Dr. Anker can be contacted at [s.anker@cachexia.de](mailto:s.anker@cachexia.de) or at the Department of Cardiology (CVK), Charité Universitätsmedizin, D-13353 Berlin, Germany.

\*A list of the RESHAPE-HF2 investigators is provided in the Supplementary Appendix, available at [NEJM.org](http://NEJM.org).

This article was published on August 31, 2024, and last updated on November 14, 2024, at [NEJM.org](http://NEJM.org).

*N Engl J Med* 2024;391:1799-809.

DOI: 10.1056/NEJMoa2314328

Copyright © 2024 Massachusetts Medical Society.

**CME**

**F**UNCTIONAL MITRAL REGURGITATION IS common among patients with heart failure and is associated with a poor prognosis.<sup>1,2</sup> Despite guideline-recommended medical therapy and cardiac-resynchronization therapy, many patients with functional mitral regurgitation remain symptomatic.<sup>3</sup> Surgery is usually not recommended unless another surgical intervention, such as coronary-artery bypass grafting or aortic-valve replacement, is indicated.<sup>4,5</sup> For patients who are not surgical candidates, current international heart-failure guidelines suggest that transcatheter edge-to-edge repair for functional mitral regurgitation should be considered, but they do not make strong recommendations<sup>6,7</sup> because of conflicting evidence of benefit. The MITRA-FR (Percutaneous Repair with the MitraClip Device for Severe Functional/Secondary Mitral Regurgitation) trial showed that percutaneous repair did not lead to a lower rate of death from any cause or hospitalization for heart failure or a lower rate of death from any cause at 1 year and 2 years than medical therapy alone.<sup>8,9</sup> In contrast, the COAPT (Cardiovascular Outcomes Assessment of the MitraClip Percutaneous Therapy for Heart Failure Patients with Functional Mitral Regurgitation) trial showed that transcatheter mitral-valve repair led to not only a lower rate of hospitalization for heart failure but also a lower rate of death from any cause than medical therapy alone during 24 months of follow-up.<sup>10</sup> Several hypotheses have been proposed to explain these discordant results, including differences in the severity of heart failure, medical treatment, and the mechanisms underlying functional mitral regurgitation.<sup>11,12</sup>

The MITRA-FR and COAPT trials involved patients with predominantly severe functional mitral regurgitation, which is usually defined by guidelines as a regurgitant volume of greater than 60 ml, a regurgitant fraction of greater than 50%, and a mean effective regurgitant orifice area larger than 0.40 cm<sup>2</sup>. The mean effective regurgitant orifice area in these trial populations were 0.31 cm<sup>2</sup> and 0.40 cm<sup>2</sup>.<sup>11,12</sup> A clinical-practice registry showed that 44% of patients undergoing transcatheter mitral-valve repair had only moderate mitral regurgitation.<sup>13</sup> Worldwide, more than 150,000 patients have undergone transcatheter mitral-valve repair, but conclusive evidence of efficacy based on the results of randomized trials is limited, especially for patients with moderate or moderate-to-severe functional mitral re-

gurgitation.<sup>14,15</sup> Accordingly, we conducted a randomized trial to provide further evidence regarding the safety and effectiveness of transcatheter mitral-valve repair in patients with symptomatic heart failure and functional mitral regurgitation.

## METHODS

### TRIAL DESIGN AND OVERSIGHT

RESHAPE-HF2 (Randomized Investigation of the MitraClip Device in Heart Failure: Second Trial in Patients with Clinically Significant Functional Mitral Regurgitation) is a prospective, multicenter, investigator-initiated randomized trial involving patients with symptomatic heart failure and moderate to severe functional mitral regurgitation (despite receiving guideline-directed medical therapy) in whom isolated mitral-valve surgery was not recommended.<sup>16-18</sup> The trial was supported by an unrestricted grant from Abbott Laboratories (the financial sponsor) to the University Medical Center Göttingen (the legal sponsor). The trial protocol (available with the full text of this article at NEJM.org) was designed by the principal investigators and representatives from Abbott Laboratories and approved by the institutional review board or ethics committee at each participating site. All the patients provided written informed consent.

An independent data and safety monitoring committee evaluated patient safety during the trial. Representatives from the Clinical Trials Unit of University Medical Center Göttingen coordinated the trial and collected the data. The first, second, and ninth authors had unrestricted access to the data and vouch for the accuracy and completeness of the data and analyses and for the fidelity of the trial to the protocol. The first draft of the manuscript was written by the first, second, and fifth authors. All statistical analyses were performed by the second and ninth authors. Earlier versions of the manuscript were reviewed and edited by all the authors.

### PATIENTS, RANDOMIZATION, AND FOLLOW-UP

Eligible patients had symptoms and signs of heart failure despite guideline-recommended therapy, grade 3+ or 4+ functional mitral regurgitation,<sup>19</sup> and a left ventricular ejection fraction of 20 to 50% (initially, left ventricular ejection fraction was 15 to 35% for patients with New York Heart Association [NYHA] functional class II heart failure and 15 to 45% for patients with NYHA functional class III or IV heart failure) and were

 A Quick Take is available at NEJM.org



required to have, within 90 days before enrollment, either a hospitalization for heart failure or an elevated plasma natriuretic peptide concentration (a B-type natriuretic peptide [BNP] level of  $\geq 300$  pg per milliliter or an N-terminal pro-BNP [NT-proBNP] level of  $\geq 1000$  pg per milliliter). Patients for whom mitral-valve surgery was recommended were not eligible. Other exclusion criteria were degenerative mitral-valve disease and any of the following procedures performed within 90 days before enrollment: percutaneous coronary intervention, cardiovascular surgery, or atrial fibrillation ablation. The severity of functional mitral regurgitation was defined according to the criteria of European Association of Echocardiography.<sup>20</sup> The complete list of inclusion and exclusion criteria is shown in Table S1 in the Supplementary Appendix, available at NEJM.org.

All the patients at each site were evaluated by a heart team consisting of a heart-failure specialist, an interventional cardiologist, an echocardiographer, and a cardiothoracic surgeon. Both transthoracic and transesophageal echocardiography were required to determine patient eligibility. Patients were considered for randomization only if their heart failure was considered to be sufficiently managed (as assessed by an investigator) and site investigators and staff at the echocardiography core laboratory had confirmed their eligibility.

Patients were randomly assigned in a 1:1 ratio to undergo transcatheter mitral-valve repair and receive medical therapy (device group) or to receive medical therapy alone (control group). Medical therapy was based on guideline recommendations directed by evidence from clinical trials. Patients in the device group were scheduled to undergo transcatheter mitral-valve repair with the Mitra-Clip device implantation within 14 days after randomization. Randomization was based on permuted blocks and was stratified according to trial site and cause of heart failure (ischemic or nonischemic). Details about the procedure have been published previously (Table S2).<sup>21,22</sup> Follow-up visits were conducted at discharge (device group only), at 30, 180, and 365 days, and then yearly. An electronic data-capture system was used to populate case-report forms. All the patients were followed up by a heart-failure specialist (Table S3).

#### TRIAL OUTCOMES

There were three primary end points: the rate of the composite of first or recurrent hospitaliza-

tion for heart failure or death from cardiovascular causes during 24 months, the rate of first or recurrent hospitalization for heart failure during 24 months, and the change from baseline to 12 months in the score on the Kansas City Cardiomyopathy Questionnaire–Overall Summary (KCCQ-OS; scores range from 0 to 100, with higher scores indicating better health status). Secondary end points included mitral regurgitation of grade 2+ or worse at 12 months, change from baseline to 12 months in the 6-minute walk distance, death from any cause during the entire trial period, the rate of first or recurrent hospitalization for any cause during 24 months, and NYHA functional class I or II heart failure at 12 months. Left ventricular volumes and severity of mitral regurgitation were assessed at an independent echocardiographic core laboratory. A complete list of trial end points is provided in Table S4.

#### STATISTICAL ANALYSIS

Assuming an annual incidence of death from cardiovascular causes of 12.5%, an annual incidence of death from noncardiovascular causes of 5.5%, and an annual incidence of hospitalization for heart failure of 60 events per 100 patient-years of follow-up in the control group and a follow-up period of 24 months, we originally estimated that a sample of 420 patients would provide the trial with 80% power to reject the null hypothesis of no effect under the assumption of a hazard ratio of 0.75 at a two-sided significance level of 5%.<sup>16</sup> In 2018, a blinded sample-size review based on noncomparative data was carried out and shared with the steering committee and the legal and financial sponsors; because blinded reviews generally do not inflate the type I error rate, no alpha adjustment was made.<sup>23</sup> After the blinded sample-size review, we decided to follow the patients for more than 24 months and enroll at least 650 patients to maintain 80% power to reject the null hypothesis at a hazard ratio of 0.70. In the aftermath of the coronavirus disease 2019 pandemic, recruitment was lower than anticipated, and ethics committees in several countries would not agree to reporting and adjudication of events after 24 months because of the wording of the original informed-consent form. Accordingly, recruitment was closed in October 2023, with a planned final follow-up visit in April 2024. On February 6, 2024, the steering committee, the members of which were unaware

of the trial-group assignments, decided to revert to the originally planned primary end point — the rate of first or recurrent hospitalization for heart failure or death from cardiovascular causes during 24 months of follow-up — and to complement it with two additional primary end points: the rate of first or recurrent hospitalization for heart failure during 24 months and the change from baseline to 12 months in the KCCQ-OS score.

Statistical analyses were performed according to the intention-to-treat principle and included all the patients who had undergone randomization. The primary and secondary end points were analyzed by means of the Hochberg procedure.<sup>24</sup> All end points that were based on recurrent events were analyzed with Lin–Wei–Yang–Ying (LWYY) models. In these analyses, trial discontinuation was handled as an independent censoring event, and the actual time in the trial was included in the LWYY analyses. Sensitivity analyses were conducted in the form of shared random-effects models that included the event of interest and competing events. Analyses of continuous outcomes, including the changes from baseline to 12 months in the KCCQ-OS score, the 6-minute walk distance, and echocardiographic variables, were performed with linear mixed models for repeated measures with gaussian distribution and included all available data with no imputation. Although the mixed models for repeated measures approach is robust with regard to missing data to some extent, sensitivity analyses were performed with the use of reference-based multiple imputation. Subgroup analyses of the first primary end point were conducted with the use of 13 baseline characteristics. The respective baseline characteristic and its interaction with treatment were included in the LWYY regression model described above. Further details are provided in the statistical analysis plan, available with the protocol. Statistical analyses were performed with the use of R software, version 4.3.1.

## RESULTS

### PATIENTS

From March 2015 through October 2023, a total of 621 patients underwent screening, of whom 505 were enrolled at 30 sites in nine countries. A total of 250 patients were randomly assigned to the device group and 255 to the control group (Fig. S1). The baseline characteristics of the patients are shown in Table 1 and Table S5. Among the trial

patients, the mean ( $\pm$ SD) age was  $70\pm 10$  years, 20% were women, 35% had nonischemic cardiomyopathy, and 29% had cardiac-resynchronization therapy devices. The median left ventricular ejection fraction was 31% (interquartile range, 25 to 37), the median left ventricular end-diastolic volume was 205 ml (interquartile range, 157 to 250), the median effective regurgitant orifice area was 0.23 cm<sup>2</sup> (interquartile range, 0.20 to 0.29), and the median regurgitant volume was 36 ml (interquartile range, 29 to 43). The median KCCQ-OS score was 43 points (interquartile range, 26 to 63). Medical therapy at baseline appeared to be similar in the two trial groups. The representativeness of the trial population is shown in Table S6.

### PROCEDURES

Of 250 participants assigned to the device group, 248 (99.2%) had transcatheter mitral-valve repair attempted. The MitraClip device was deployed in 244 patients (98.4%), and echocardiographic data at the end of the procedure were available for 243 patients (97.2%). The severity grade of mitral regurgitation was 1+ or lower in 181 patients (74.5%), 2+ in 43 patients (17.7%), 3+ in 10 patients (4.1%), and 4+ in 9 patients (3.7%). Procedures were performed within 14 days after randomization in 220 patients, within 15 to 30 days in 21 patients, and after 30 days in 7 patients. The median procedure time was 117 minutes (interquartile range, 70 to 150). The median fluoroscopy time was 22 minutes (interquartile range, 12 to 34).

### EFFICACY END POINTS

The mean duration of follow-up was  $18.8\pm 8.2$  months. At 24 months, the rate of first or recurrent hospitalization for heart failure or death from cardiovascular causes was 37.0 events per 100 patient-years (a total of 151 events) in the device group and 58.9 events per 100 patient-years (a total of 225 events) in the control group (rate ratio, 0.64; 95% confidence interval [CI], 0.48 to 0.85;  $P=0.002$ ). The rate of first or recurrent hospitalization for heart failure during 24 months was 26.9 events per 100 patient-years (a total of 110 events) in the device group and 46.6 events per 100 patient-years (a total of 178 events) in the control group (rate ratio, 0.59; 95% CI, 0.42 to 0.82;  $P=0.002$ ).

The mean change from baseline to 12 months in the KCCQ-OS score was  $21.6\pm 26.9$  points in the device group and  $8.0\pm 24.5$  points in the control group (least-squares mean difference, 10.9 points;

**Table 1. Baseline Demographic and Clinical Characteristics of the Patients with Heart Failure and Moderate to Severe Functional Mitral Regurgitation.\***

Characteristic	Device Group (N=250)	Control Group (N=255)
Age — yr	70.0±10.4	69.4±10.7
Male sex — no. (%)	195 (78.0)	211 (82.8)
Diabetes — no. (%)	91 (36.4)	85 (33.3)
Hypertension — no. (%)	141 (56.4)	127 (49.8)
Previous myocardial infarction — no. (%)	144 (57.6)	135 (52.9)
Previous PCI — no. (%)	118 (47.2)	125 (49.0)
Previous CABG — no. (%)	69 (27.6)	64 (25.1)
Previous stroke or TIA — no. (%)	29 (11.6)	30 (11.8)
Peripheral vascular disease — no. (%)	38 (15.2)	27 (10.6)
History of atrial fibrillation or flutter — no. (%)	118 (47.2)	125 (49.0)
Body-mass index†	27.0±4.3	26.7±4.3
Nonischemic cause of cardiomyopathy — no. (%)	88 (35.2)	88 (34.5)
NYHA functional class — no. (%)		
II	59 (23.6)	65 (25.5)
III	150 (60.0)	153 (60.0)
IV	41 (16.4)	36 (14.1)
Hospitalization for heart failure within previous year — no. (%)	165 (66.0)	168 (65.9)
Systolic blood pressure — mm Hg	112±16	113±16
Median NT-proBNP level (IQR) — pg/ml‡	2651 (1630–4918)	2816 (1306–5496)
Median BNP level (IQR) — pg/ml§	556 (312–1018)	406 (231–874)
Median 6-minute walk distance (IQR) — m	300 (220–382)	310 (200–378)
Estimated glomerular filtration rate — ml/min/1.72 cm <sup>2</sup>	54.9±19.0	56.7±23.3
Median KCCQ-OS score (IQR) — points¶	42.2 (28.3–62.0)	44.3 (25.8–64.2)
Median left ventricular ejection fraction (IQR) — %	32 (26–37)	31 (25–37)
Median left ventricular end-diastolic volume (IQR) — ml	200 (153–249)	206 (158–250)
Severity of mitral regurgitation — no. (%)		
Grade 3+	141 (56.4)	141 (55.3)
Grade 4+	109 (43.6)	114 (44.7)
Median effective regurgitant orifice area (IQR) — cm <sup>2</sup>	0.23 (0.20–0.30)	0.23 (0.19–0.29)
Median regurgitant volume (IQR) — ml	35.4 (28.9–43.9)	35.6 (28.2–42.5)

\* Plus–minus values are means ±SD. CABG denotes coronary-artery bypass graft, IQR interquartile range, NYHA New York Heart Association, PCI percutaneous coronary intervention, and TIA transient ischemic attack.

† The body-mass index is the weight in kilograms divided by the square of the height in meters.

‡ Data on N-terminal pro-B-type natriuretic peptide (NT-proBNP) level were available for 384 patients (191 in the device group and 193 in the control group).

§ Data on B-type natriuretic peptide (BNP) level were available for 123 patients (61 in the device group and 62 in the control group).

¶ Scores on the Kansas City Cardiomyopathy Questionnaire–Overall Summary (KCCQ-OS) range from 0 to 100, with higher scores indicating better health status.

95% CI, 6.8 to 15.0; P<0.001). The results for the primary and secondary end points are shown in Table 2 and Figure 1. The total number of first or recurrent hospitalizations for any cause during 24 months was 199 in the device group and 233

in the control group (rate ratio, 0.82; 95% CI, 0.63 to 1.07). During the entire duration of follow-up (38.1±18.1 months), 142 deaths occurred in the device group (annualized rate [the number of events divided by the total number of patient-

**Table 2. Primary and Secondary End Points.\***

End Point	Device Group (N=250)	Control Group (N=255)	Difference (95% CI)†	P Value‡
<b>Primary End Points§</b>				
Rate of the composite of first or recurrent hospitalization for heart failure or death from cardiovascular causes during 24 mo per 100 patient-yr (no. of events/total no. of patient-yr)	37.0 (151/408.6)	58.9 (225/381.9)	0.64 (0.48–0.85)	0.002¶
Rate of first or recurrent hospitalization for heart failure during 24 mo per 100 patient-yr (no. of events/total no. of patient-yr)	26.9 (110/408.6)	46.6 (178/381.9)	0.59 (0.42–0.82)	0.002¶
Mean change from baseline to 12 mo in the KCCQ-OS score — points	21.6±26.9	8.0±24.5	10.9 (6.8–15.0)‖	<0.001¶
<b>Secondary end points</b>				
Mitral regurgitation grade ≤2+ at 12 mo — no./total no. (%)	132/146 (90.4)	43/119 (36.1)**	21.3 (10.7–45.8)††	<0.001¶
Mean change in 6-min walk distance from baseline to 12 mo — m	34.0±105.9	5.1±97.6	20.5 (0.3–40.7)‖	0.05‡‡
Rate of death from any cause during the complete follow-up per 100 patient-yr (no. of events/total no. of patient-yr)	17.0 (142/836.7)	18.6 (142/765.2)	0.90 (0.71–1.13)	0.37‡‡
Rate of first or recurrent hospitalization for any cause during 24 mo per 100 patient-yr (no. of events/total no. of patient-yr)	48.7 (199/408.6)	61.0 (233/381.9)	0.82 (0.63–1.07)	0.15‡‡
NYHA functional class I or II heart failure at 12 mo — no./total no. (%)§§	140/188 (74.5)	96/164 (58.5)	2.35 (1.48–3.77)††	<0.001¶

\* Plus-minus values are means ±SD.

† The difference in treatment effect is given as a rate ratio unless otherwise noted.

‡ Unadjusted P values are presented, but they were tested for significance according to the thresholds as defined by the Hochberg procedure.

§ The three primary end points were tested at a two-sided significance level of 5%. After the results for all three were found to be statistically significant, the Hochberg procedure was also applied to the secondary end points at a two-sided significance level of 5%.

¶ The finding was statistically significant according to the Hochberg procedure.

‖ The difference in treatment effect is given as the difference in least-squares means, as estimated with linear mixed models for repeated measures with gaussian distribution.

\*\* Of these 43 patients, 15 had undergone transcatheter mitral-valve repair.

†† The difference in treatment effect is given as an odds ratio.

‡‡ The finding was not significant according to the Hochberg procedure.

§§ At 12 months, 33 patients in the control group and 27 patients in the device group had died.

years), 17.0%) and 142 deaths occurred in the control group (annualized rate, 18.6%), for a hazard ratio of 0.90 (95% CI, 0.71 to 1.13). A total of 140 of 188 patients (74.5%) in the device group and 96 of 164 (58.5%) in the control group had NYHA functional class I or II heart failure at 12 months ( $P<0.001$ ). The mean change in 6-minute walk distance from baseline to 12 months was 34.0±105.9 m in the device group and 5.1±97.6 m in the control group (least-squares mean difference, 20.5 m; 95% CI, 0.3 to 40.7;  $P=0.047$ ; not significant according to the Hochberg procedure). A total of 132 of 146 patients (90.4%) in the device group and 43 of 119 patients (36.1%) in the control group had a mitral regurgitation severity grade of 2+ or lower at 12 months ( $P<0.001$ ). The results of the subgroup analyses are shown in Figure 2.

## SAFETY

Adverse events are shown in Table 3. Among the patients assigned to the device group, periprocedural adverse events were reported in four (1.6%). These events included two cases of hematoma, one case of pericardial effusion, and one case of right atrial perforation that led to thoracotomy after completion of the device implantation.

## DISCUSSION

In the RESHAPE-HF2 trial, which involved patients with moderate to severe functional mitral regurgitation who remained symptomatic despite medical therapy, transcatheter mitral-valve repair led to a lower rate of first or recurrent hospitalization for heart failure or death from cardiovascular


**Figure 1. Number of Primary End-Point Events and Change in KCCQ-OS Score.**


Panel A shows the total number of events in the composite of first or recurrent hospitalization for heart failure or death from cardiovascular causes over a period of 24 months. Panel B shows the total number of first or recurrent hospitalizations for heart failure over a period of 24 months. Panel C shows the least-squares mean change from baseline over a period of 12 months in the score on the Kansas City Cardiomyopathy Questionnaire—Overall Summary (KCCQ-OS) for quality of life (scores range from 0 to 100, with higher scores indicating better health status). I bars indicate 95% confidence intervals.

causes during 24 months, as well as a lower rate of first or recurrent hospitalization for heart failure during 24 months and a greater increase from baseline to 12 months in the KCCQ-OS score, than medical therapy alone. These results appeared to be consistent across most subgroups.

Previous studies of transcatheter mitral-valve repair in patients with functional mitral regurgitation mainly involved patients with severe mitral regurgitation and provided conflicting results.<sup>8,10</sup> The extent of left ventricular damage and severity of mitral-valve regurgitation have been proposed as important determinants of benefit from transcatheter mitral-valve repair.<sup>25,26</sup> The mean effective regurgitant orifice area was smaller among the patients in our trial (0.25 cm<sup>2</sup>) than among those in the COAPT trial (0.40 cm<sup>2</sup>) and the MITRA-FR trial (0.31 cm<sup>2</sup>); at baseline, 14% of the patients in our trial had an effective regurgitant orifice area larger than 0.40 cm<sup>2</sup> and 23% had an effective regurgitant orifice area smaller than 0.20 cm<sup>2</sup>. However, there was no apparent heterogeneity in treatment effect in subgroups defined according to key baseline echocardiographic measurements. The findings of the current trial are relevant because untreated functional mitral regurgitation can cause changes in cardiac structure and function that lead to worsening heart failure and an adverse prognosis.<sup>27,28</sup>

Patients with heart failure and functional mitral regurgitation are at high risk for recurrent hospitalizations.<sup>1,29</sup> The relative difference in the rate of first or recurrent hospitalization for heart failure during 24 months between the device group and control group of 41% represents an absolute difference in risk of 19.7 events per





**Figure 2. Subgroup Analysis of the Composite of First or Recurrent Hospitalization for Heart Failure or Death from Cardiovascular Causes.**

The annualized rate (expressed as a percentage) is the number of events divided by the total number of patient-years. The body-mass index is the weight in kilograms divided by the square of the height in meters. BNP denotes B-type natriuretic peptide level, LVEF left ventricular ejection fraction, NT-proBNP N-terminal pro-BNP, and NYHA New York Heart Association.

100 patient-years, which translates into a number needed to treat of only 5.1 to prevent one hospitalization for heart failure with transcatheter mitral-valve repair in combination with guideline-directed

medical therapy. Also, the reductions in symptoms that were assessed according to NYHA functional class and the KCCQ-OS score at 12 months were greater in the device group than in the control

**Table 3. Adverse Events of Interest during 24 Months in the Intention-to-Treat Population.\***

Event	Device Group (N=250)	Control Group (N=255)	Hazard or Rate Ratio (95% CI)†	P Value
<i>no. of patients with event (estimate of event rate)</i>				
Death from any cause‡	51 (22.3)	67 (29.6)	0.73 (0.51–1.05)	0.09
Death from cardiovascular causes§	41 (17.8)	47 (20.4)	0.84 (0.55–1.28)	0.43
Death from noncardiovascular causes§	10 (4.5)	20 (9.3)	0.46 (0.22–0.99)	0.04
Unplanned MitraClip implantation¶	8 (2.0)	25 (6.5)	0.32 (0.14–0.70)	0.004
All unplanned transcatheter mitral-valve repair¶	8 (2.0)	38 (10.0)**	0.21 (0.10–0.44)	<0.001
Mitral-valve surgery††	1 (0.004)	2 (0.008)	0.51 (0.05–5.58)	0.57
PCI§	6 (0.026)	8 (0.034)	0.74 (0.26–2.12)	0.57
CABG	0	0	—	—
Stroke§	5 (0.022)	2 (0.008)	2.5 (0.48–12.9)	0.25
Myocardial infarction¶	3 (0.007)	3 (0.008)	1.02 (0.14–7.52)	0.99
LVAD implantation††	1 (0.008)	2 (0.02)	0.5 (0.05–5.49)	0.56
Heart transplantation	1	0	—	—
Implantation of ICD or CRT-D§	7 (1.8)	7 (1.7)	0.96 (0.35–2.66)	0.93

\* CRT-D denotes cardiac-resynchronization therapy defibrillator, and ICD implantable cardioverter-defibrillator.  
 † Whenever recurrent-event analyses were performed (unplanned MitraClip implantation, all unplanned transcatheter mitral-valve repair, and myocardial infarction), rate ratios are given.  
 ‡ Death from any cause was analyzed by means of Cox regression, with adjustment for country and cause of heart failure. Event rates are given as Kaplan–Meier estimates.  
 § Death from cardiovascular causes, death from noncardiovascular causes, PCI, stroke, and implantation of ICD or CRT-D were analyzed by means of Cox regression, with adjustment for country and cause of heart failure. Event rates are given as the cumulative incidence (expressed as a percentage) at 24 months. In these analyses, death was included in the model as a competing event.  
 ¶ The analyses of unplanned MitraClip implantations, all unplanned transcatheter mitral-valve repair, and myocardial infarction included recurrent events and were conducted with the Lin–Wei–Yang–Ying model for recurrent events, with adjustment for country and cause of heart failure. Event rates are given as the annualized rate, defined as the number of events divided by the total number of patient-years and expressed as a percentage.  
 || All eight implantations were performed by month 12 (with seven performed by month 6).  
 \*\* Of the 38 implantations, 35 were performed by month 12 (with 28 performed by month 6).  
 †† Mitral-valve surgery and left ventricular assist device (LVAD) implantation were analyzed by means of Cox regression without covariate adjustments. Event rates are given as the cumulative incidence (expressed as a percentage) at 24 months. In the analyses, death was included in the model as a competing event.

group, and the difference between the groups in the 6-minute walk distance at 12 months was a least-squares mean of 20.5 m in favor of the device group. By 12 months, 90.4% (132 of 146) of the patients assigned to the device group had a reduction in mitral regurgitation grade to 2+ or lower, as compared with 36% (43 of 119) of those in the control group, of whom 35% (15 of 43) had undergone a mitral-valve procedure. This finding may explain the greater reductions in symptoms and hospitalization rates and improvements in functional capacity and health status that were observed in device group as compared with the control group.

The COAPT trial showed that during 24 months of follow-up, transcatheter mitral-valve repair in combination with medical therapy resulted in a lower rate of death from any cause than medical

therapy alone (hazard ratio, 0.62; 95% CI, 0.46 to 0.82),<sup>10</sup> but the MITRA-FR trial did not show a lower rate of death from any cause with transcatheter mitral-valve repair in a similar comparison.<sup>8</sup> In our trial, the rate of death from any cause was also not lower with transcatheter mitral-valve repair than with medical therapy alone, which could be related to a smaller clinical benefit or fewer outcome events, findings consistent with both less severe mitral regurgitation and less advanced heart failure. All-cause mortality at 2 years in the control group of our trial was 29.6% (Table 3), as compared with 46.1% in control group of the COAPT trial, and the rate of first or recurrent hospitalization in our trial was less than half that observed in the COAPT trial, which suggests that the patients enrolled in our trial had less severe disease.

The trial has several limitations. Although randomization was performed in a blinded manner, the participants, investigators, and echocardiographers were not unaware of subsequent treatments. This situation could have created bias, especially for quality-of-life assessments recorded by patients. Some patients who had been assigned to receive medical therapy alone underwent transcatheter mitral-valve repair, which could have diluted the observed treatment effect. Finally, the trial was not designed to show differences in mortality.

Among patients with heart failure and moderate to severe functional mitral regurgitation who remained symptomatic despite guideline-recommended medical therapy, transcatheter mitral-valve repair in combination with medical therapy

led to a lower rate of first or recurrent hospitalization for heart failure or death from cardiovascular causes during 24 months, a lower rate of first or recurrent hospitalization for heart failure during 24 months, and better health status at 12 months than medical therapy alone.

Supported by an unrestricted grant of Abbott Laboratoriosto University Medical Center Göttingen. The development of the Web-based end-point assessment tool that was used in the event adjudication process in this trial was supported by a grant from Stiftung Charité.

Disclosure forms provided by the authors are available with the full text of this article at NEJM.org.

A data sharing statement provided by the authors is available with the full text of this article at NEJM.org.

We thank Volker Möller (project leader in the Center for Cardiovascular Telemedicine of Charité Universitätsmedizin Berlin) for assistance in setting up the end-point assessment tool for use in the adjudication process.

#### APPENDIX

The authors' full names and academic degrees are as follows: Stefan D. Anker, M.D., Tim Friede, Ph.D., Ralph-Stephan von Bardeleben, M.D., Javed Butler, M.D., Muhammad-Shahzeb Khan, M.D., Monika Diek, M.Sc., Jutta Heinrich, M.Sc., Martin Geyer, M.D., Marius Placzek, Ph.D., Roberto Ferrari, M.D., William T. Abraham, M.D., Ottavio Alfieri, M.D., Angelo Auricchio, M.D., Antoni Bayes-Genis, M.D., John G.F. Cleland, M.D., Gerasimos Filippatos, M.D., Finn Gustafsson, M.D., Wilhelm Haverkamp, M.D., Malte Kelm, M.D., Karl-Heinz Kuck, M.D., Ulf Landmesser, M.D., Aldo P. Maggioni, M.D., Marco Metra, M.D., Vlasios Ninios, M.D., Mark C. Petrie, M.D., Tienush Rassaf, M.D., Frank Ruschitzka, M.D., Ulrich Schäfer, M.D., P. Christian Schulze, M.D., Konstantinos Spargias, M.D., Alec Vahanian, M.D., Jose Luis Zamorano, M.D., Andreas Zeiher, M.D., Mahir Karakas, M.D., Friedrich Koehler, M.D., Miĳa Lainscak, M.D., Alper Öner, M.D., Nikolaos Mezilis, M.D., Efstratios K. Theofilogiannakos, M.D., Ilias Ninios, M.D., Michael Chrissoheris, M.D., Panagiota Kourkovi, M.D., Konstantinos Papadopoulos, M.D., Grzegorz Smolka, M.D., Wojciech Wojakowski, M.D., Krzysztof Reczuch, M.D., Fausto J. Pinto, M.D., Łukasz Wiewiórka, M.D., Zbigniew Kalarus, M.D., Marianna Adamo, M.D., Evelyn Santiago-Vacas, M.D., Tobias F. Ruf, M.D., Michael Gross, M.D., Joern Tongers, M.D., Gerd Hasenfuss, M.D., Wolfgang Schillinger, M.D., and Piotr Ponikowski, M.D.

The authors' affiliations are as follows: the Department of Cardiology of German Heart Center Charité and the Institute of Health Center for Regenerative Therapies, German Center for Cardiovascular Research (DZHK) Partner Site Berlin, Charité Universitätsmedizin (S.D.A., M.D., W.H.), the Department of Cardiology, Angiology, and Intensive Care Medicine, Deutsches Herzzentrum der Charité, Campus Benjamin Franklin, the Berlin Institute of Health, and DZHK Partner Site Berlin (U.L.), Deutsches Herzzentrum der Charité, the Department of Cardiology, Angiology, and Intensive Care Medicine, Campus Charité Mitte, the Center for Cardiovascular Telemedicine, and DZHK Partner Site Berlin, Charité Universitätsmedizin (F.K.), Berlin, the Departments of Cardiology and Pneumology (S.D.A., M.D., G.H., W.S.) and Medical Statistics (T.F., M.P.) and the Clinical Trial Unit (J.H.), University Medical Center Göttingen, and the Heart Center, Department of Cardiology (W.S.), Georg August University of Göttingen, DZHK Partner Site Göttingen, Göttingen, the Department of Cardiology, University Medical Center of the Johannes Gutenberg–University Mainz (R.S.B., M. Geyer, T.F.R.), Mainz, the Department of Cardiology, Pulmonology, and Vascular Medicine, University Hospital Düsseldorf, Medical Faculty of the Heinrich Heine University Düsseldorf, and Cardiovascular Research Institute Düsseldorf, Medical Faculty and University Hospital Düsseldorf, Heinrich Heine University, Düsseldorf (M. Kelm), the Department of Rhythmology, University Heart Center Lübeck, University Hospital Schleswig–Holstein, Lübeck (K.-H.K.), the West German Heart and Vascular Center, Department of Cardiology and Vascular Medicine, University Hospital Essen, University Duisburg–Essen, Essen (T.R.), the Heart and Vascular Center, Bad Bevensen (U.S.), the Department of Internal Medicine I, Cardiology, University Hospital Jena, Jena (P.C.S.), the Institute for Cardiovascular Regeneration, Goethe University Frankfurt, and DZHK Partner Site Rhein–Main, Frankfurt am Main (A.Z.), the Department of Intensive Care Medicine, University Medical Center Hamburg–Eppendorf, and DZHK Partner Site Hamburg–Kiel–Lübeck, Hamburg (M. Karakas), the Department of Cardiology, Rostock University Medical Center, Rostock (A.Ö.), the Department of Medicine, Heart and Vascular Center, Division of Cardiology and Vascular Medicine, Johanniter Hospital Stendal, Stendal (M. Gross), the Mid-German Heart Center, Department of Cardiology, Angiology, and Intensive Care Medicine, University Hospital Halle, Halle (J.T.) — all in Germany; the Department of Medicine, University of Mississippi Medical Center, Jackson (J.B.); Baylor Scott and White Research Institute, Dallas (J.B., M.S.K.), Baylor Scott and White the Heart Hospital Plano, Plano (M.S.K.), and the Department of Medicine, Baylor College of Medicine, Temple (M.S.K.) — all in Texas; the Department of Translational Medicine, University of Ferrara, Ferrara (R.F.), the Cardiac Surgery Unit, IRCCS San Raffaele Scientific Institute, Milan (O.A.), ANMCO Associazione Nazionale Medici Cardiologi Ospedalieri Research Center, Heart Care Foundation, Florence (A.P.M.), and the Institute of Cardiology, Azienda Socio Sanitaria Territoriale degli Spedali Civili di Brescia, and the Department of Medical and Surgical Specialties, Radiologic Sciences and Public Health, University of Brescia, Brescia (M.M., M.A.) — all in Italy; the Division of Cardiovascular Medicine, Ohio State University, Columbus (W.T.A.); the Department of Cardiology, Cardiocentro Ticino Institute, Ente Ospedaliero Cantonale, Lugano (A.A.), and the Center for Translational and Experimental Cardiology, Department of Cardiology, University Hospital Zurich, University of Zurich, Zurich (F.R.) — both in Switzerland; the Heart Institute, Hospital Universitari Germans Trias i Pujol, Barcelona (A.B.-G., E.S.-V.), and the Department of Cardiology, Hospital Universitario Ramón y Cajal, and Centro de Investigación Biomédica en Red de Enfermedades Cardiovasculares, Instituto de Salud Carlos III, Madrid (J.L.Z.) — all in Spain; the School of Cardiovascular and Metabolic Health, University of Glasgow, British Heart Foundation Glasgow Cardiovascular Research Centre, Glasgow, United Kingdom (J.G.F.C., M.C.P.); the Department of Cardiology, Attikon University Hospital, School of Medicine, National and Kapodistrian University of Athens (G.F.), and the Department of Transcatheter Heart

Valves, Hygeia Hospital (K.S., M.C., P.K., K.P.), Athens, and the Department of Cardiology, St. Luke's Hospital (N.M., E.K.T.), and the European Interbalkan Medical Center (V.N., I.N., K.P.), Thessaloniki — all in Greece; the Department of Cardiology, Copenhagen University Hospital—Rigshospitalet, Copenhagen (F.G.); Unité Formation et Recherche Médecine, Université de Paris-Cité, site Bichat, Laboratoire de Recherche Vasculaire Translationnelle, INSERM, Groupe Hospitalier Bichat, Paris (A.V.); the Division of Cardiology, General Hospital Murska Sobota, Murska Sobota, and the Faculty of Medicine, University of Ljubljana, Ljubljana — both in Slovenia (M.L.); the Division of Cardiology and Structural Heart Diseases (G.S., W.W.) and the Department of Cardiology, Congenital Heart Diseases, and Electrotherapy (Z.K.), Medical University of Silesia, Katowice, the Institute of Heart Diseases, Medical University and University Hospital, Wrocław (K.R., P.P.), and the Department of Interventional Cardiology, Department of Radiology and Diagnostic Imaging, John Paul II Hospital, Krakow (Ł.W.) — all in Poland; and Centro Academico de Medicina de Lisboa, Faculdade de Medicina da Universidade de Lisboa, Lisbon, Portugal (F.J.P.).

## REFERENCES

- Giustino G, Camaj A, Kapadia SR, et al. Hospitalizations and mortality in patients with secondary mitral regurgitation and heart failure: the COAPT trial. *J Am Coll Cardiol* 2022;80:1857-68.
- Adamo M, Chioncel O, Benson L, et al. Prevalence, clinical characteristics and outcomes of heart failure patients with or without isolated or combined mitral and tricuspid regurgitation: an analysis from the ESC-HFA Heart Failure Long-Term Registry. *Eur J Heart Fail* 2023;25:1061-71.
- Barnes S, Sharma H, Gamble J, Dawkins S. Management of secondary mitral regurgitation: from drugs to devices. *Heart* 2023 August 22 (Epub ahead of print).
- Verma S, Mesana TG. Mitral-valve repair for mitral-valve prolapse. *N Engl J Med* 2009;361:2261-9.
- Nishimura RA, Otto CM, Bonow RO, et al. 2017 AHA/ACC focused update of the 2014 AHA/ACC guideline for the management of patients with valvular heart disease: a report of the American College of Cardiology/American Heart Association Task Force on Clinical Practice Guidelines. *J Am Coll Cardiol* 2017;70:252-89.
- McDonagh TA, Metra M, Adamo M, et al. 2023 Focused update of the 2021 ESC guidelines for the diagnosis and treatment of acute and chronic heart failure: developed by the task force for the diagnosis and treatment of acute and chronic heart failure of the European Society of Cardiology (ESC) with the special contribution of the Heart Failure Association (HFA) of the ESC. *Eur J Heart Fail* 2024;26:5-17.
- Heidenreich PA, Bozkurt B, Aguilar D, et al. 2022 AHA/ACC/HFSA guideline for the management of heart failure: executive summary: a report of the American College of Cardiology/American Heart Association Joint Committee on Clinical Practice Guidelines. *J Am Coll Cardiol* 2022;79:1757-80.
- Obadia J-F, Messika-Zeitoun D, Leurent G, et al. Percutaneous repair or medical treatment for secondary mitral regurgitation. *N Engl J Med* 2018;379:2297-306.
- Iung B, Armoiry X, Vahanian A, et al. Percutaneous repair or medical treatment for secondary mitral regurgitation: outcomes at 2 years. *Eur J Heart Fail* 2019;21:1619-27.
- Stone GW, Lindenfeld J, Abraham WT, et al. Transcatheter mitral-valve repair in patients with heart failure. *N Engl J Med* 2018;379:2307-18.
- Grayburn PA, Sannino A, Packer M. Proportionate and disproportionate functional mitral regurgitation: a new conceptual framework that reconciles the results of the MITRA-FR and COAPT trials. *JACC Cardiovasc Imaging* 2019;12:353-62.
- Pibarot P, Delgado V, Bax JJ. MITRA-FR vs. COAPT: lessons from two trials with diametrically opposed results. *Eur Heart J Cardiovasc Imaging* 2019;20:620-4.
- Kar S, von Bardeleben RS, Rottbauer W, et al. Contemporary outcomes following transcatheter edge-to-edge repair: 1-year results from the EXPAND study. *JACC Cardiovasc Interv* 2023;16:589-602.
- Hibbert B, Alkhouli M. Optimal outcomes in real-world patients treated with TEER: reaching the pinnacle. *JACC Cardiovasc Interv* 2023;16:603-5.
- Mack M, Carroll JD, Thourani V, et al. Transcatheter mitral valve therapy in the United States: a report from the STS-ACC TVT Registry. *J Am Coll Cardiol* 2021;78:2326-53.
- Anker SD, Friede T, von Bardeleben RS, et al. Randomized investigation of the MitraClip device in heart failure: design and rationale of the RESHAPE-HF2 trial design. *Eur J Heart Fail* 2024;26:984-93.
- Anker SD, Friede T, von Bardeleben RS, et al. Percutaneous repair of moderate-to-severe or severe functional mitral regurgitation in patients with symptomatic heart failure: baseline characteristics of patients in the RESHAPE-HF2 trial and comparison to COAPT and MITRA-FR trials. *Eur J Heart Fail* 2024;26:1608-15.
- von Bardeleben RS, Khan MS, Geyer M, et al. Baseline echocardiographic characteristics of patients enrolled in the randomized investigation of MitraClip device in heart failure (RESHAPE HF-2) trial: comparison to COAPT and Mitra-FR. *Glob Cardiol* 2024;2:67-74 (<https://www.globalcardiology.info/site/article/view/40>).
- Stone GW, Vahanian AS, Adams DH, et al. Clinical trial design principles and endpoint definitions for transcatheter mitral valve repair and replacement: part 1: clinical trial design principles: a consensus document from the mitral valve academic research consortium. *J Am Coll Cardiol* 2015;66:278-307.
- Lancellotti P, Pibarot P, Chambers J, et al. Recommendations for the imaging assessment of prosthetic heart valves: a report from the European Association of Cardiovascular Imaging endorsed by the Chinese Society of Echocardiography, the Inter-American Society of Echocardiography, and the Brazilian Department of Cardiovascular Imaging. *Eur Heart J Cardiovasc Imaging* 2016;17:589-90.
- Sherif MA, Paranskaya L, Yuecel S, et al. MitraClip step by step: how to simplify the procedure. *Neth Heart J* 2017;25:125-30.
- Feldman T, Foster E, Glower DD, et al. Percutaneous repair or surgery for mitral regurgitation. *N Engl J Med* 2011;364:1395-406.
- Food and Drug Administration. Adaptive design clinical trials for drugs and biologics. Guidance for industry. December 2019 (<https://www.fda.gov/regulatory-information/search-fda-guidance-documents/adaptive-design-clinical-trials-drugs-and-biologics-guidance-industry>).
- Anker SD, Butler J, Talha KM, Friede T. Using multiple primary endpoints in clinical trials with a focus on heart failure. *Glob Cardiol* 2024;2:11-6 (<https://www.globalcardiology.info/site/article/view/33>).
- Stolz L, Doldi PM, Sannino A, Hausleiter J, Grayburn PA. The evolving concept of secondary mitral regurgitation phenotypes: lessons from the M-TEER trials. *JACC Cardiovasc Imaging* 2024;17:659-68.
- Packer M, Grayburn PA. Contrasting effects of pharmacological, procedural, and surgical interventions on proportionate and disproportionate functional mitral regurgitation in chronic heart failure. *Circulation* 2019;140:779-89.
- Stolz L, Doldi PM, Orban M, et al. Staging heart failure patients with secondary mitral regurgitation undergoing transcatheter edge-to-edge repair. *JACC Cardiovasc Interv* 2023;16:140-51.
- Singh GK, Namazi F, Hirasawa K, et al. Extramitral valvular cardiac involvement in patients with significant secondary mitral regurgitation. *Am J Cardiol* 2022;162:143-9.
- Stone GW, Abraham WT, Lindenfeld J, et al. Five-year follow-up after transcatheter repair of secondary mitral regurgitation. *N Engl J Med* 2023;388:2037-48.

Copyright © 2024 Massachusetts Medical Society.